## Healthy Vermonters 2020 - Windham County Quick Reference

Stanting Management and 2020 Indicates	County	County		U.S.	
Healthy Vermonters 2020 Indicator	Baseline	Current	VT Current	Current	Tar
CCESS TO HEALTH SERVICES	070/ /207	040//02:- :=:	0.404 /== :=:	2	
ercent of adults age 18-64 with health insurance	87% (2008-12)	91% (2011-15)	94% (2015)	87%	
ercent of children age 17 or younger with health insurance	98% (2008-12)	99% (2011-15)	99% (2015)	95%	
imary Care Provider FTEs per 100,000 Vermonters - Physicians (MD and DO)	94.7 (2010)	86.3 (2014)	76.1 (2014)	NA	
imary Care Provider FTEs per 100,000 Vermonters - Physician Assistants	10.8 (2010)	11.2 (2016)	13.1 (2016)	NA	
imary Care Provider FTEs per 100,000 Vermonters - Nurse Practitioners	NA	35.8 (2015)	28.7 (2015)	NA	
imary Care Provider FTEs per 100,000 Vermonters - Dentists	41.0 (2009)	43.2 (2015)	38.3 (2015)	NA	
rcent of adults with a usual primary care provider	88% (2011-12)	89% (2015-16)	88% (2016)	78%	
rcent of adults who cannot obtain care or delay care	14% (2011-12)	7% (2015-16)	8% (2016)	13%	
RTHRITIS & OSTEOPOROSIS ercent of adults with diagnosed arthritis who have activity limitations	54% (2011,13)	56% (2013,15)	49% (2015)	50%	
ercent of adults with diagnosed arthritis who engage in leisure time physical activity	83% (2011-12)	75% (2015-16)	77% (2016)		
NCER	0370 (2011 12)	7570 (2013-10)	7770 (2010)	0570 (2015)	
rerall cancer death rate per 100,000 Vermonters	169.9 (2008-10)	181.6 (2011-13)	165.9 (2013)	163.2	
rcent of adult cancer survivors always or usually getting emotional support	80% (2008, 10)	75% (2014,16)	82% (2016)	81% (2015)	
rcent of adult cancer survivors who report good to excellent general health	72% (2011,12)	71% (2015-16)	68% (2016)	65% (2015)	
rcent of woman age 21-65 receiving cervical cancer screening	89% (2006,08,10)	86% (2012,14)	86% (2014)	82%	
rcent of adults age 50-75 receiving colorectal cancer screening	66% (2008,10)	64% (2014,16)	72% (2016)	67%	
creent of female adults age 50-74 receiving breast cancer screening	80% (2006,08,10)	80% (2014,16)	79% (2016)	78%	
rcent of adolescents in grades 9-12 who reported having a sunburn in the past 12 months	NA	65% (2015)	65% (2015)	56%	
ercent of addresseries in grades 5-12 who reported having a sunburn in the past 12 months	NA	57% (2015)	54% (2015)	NA	
ite of obesity-related cancers per 100,000 Vermonters		215.8 (2011-13)	194.4 (2013)	NA.	
ite of obesity-related cancers per 100,000 Vermonters		195.8 (2011-13)	177.8 (2013)	NA NA	
ABETES & CHRONIC KIDNEY DISEASE	199.8 (2008-10)	193.8 (2011-13)	177.8 (2013)	INA	
ate of new cases of end-stage renal disease (ESRD) per million population	202.5 (2008-10)	114.5 (2013-15)	191.7 (2015)	379.9	
ercent of adults with diagnosed diabetes with A1C > 9%	NA	NA	NA	NA	
ercent of adults with diagnosed diabetes who had an annual dilated eye exam	57% (2008-10)	50% (2012,15)	67% (2015)	61%	
ercent of adults with diagnosed diabetes who had diabetes education	35% (2008-10)	19% (2012,15)	46% (2015)	55%	
ercent of adults with diagnosed prediabetes who had diabetes education ercent of adults with diagnosed prediabetes meeting aerobic physical activity guidelines	33% (2006-10) NA	19% (2012,15) NA	50% (2013)		
ARLY CHILDHOOD SCREENING	IVA	IVA	JU/0 (ZUIJ)	.5% (2013)	
ercent of children who have a developmental screening in the first 3 years of life	NA	NA	58% (2016)	NA	
ercent of newborns screened for hearing loss by 1 month of age	96% (2010)	95% (2013)	97% (2015)	83% (2012)	
ercent of newborns not passing screening, who have an audiologic evaluation by 3 months of age	NA	NA	64% (2015)	NC	
ercent of infants with hearing loss who receive intervention services by 6 months of age	NA	NA	67% (2015)		
ercent of illumina with hearing loss who receive liner vention services by 6 months or age	NA NA	NA NA	100% (2015)	NC	
IVIRONMENTAL HEALTH & FOOD SAFETY	NA NA	INA.	100% (2013)	IVC	
ercent of Vermonters served by public community water supplies that meet Safe Drinking Water Act standards	NA	NA	100% (2017)	93% (2011)	
ercent of children age 1-5 who have elevated blood lead levels (>10 μg/dL)	1.2% (2008-10)	1.2% (2013-15)	0.5% (2016)	0.50%	
ercent of children age 1-5 who have elevated blood lead levels (5-9 µg/dL)	2.3% (2010-12)	•	1.4% (2016)		
evated blood lead level (>10 µg/dL) from work exposure rate per 100,000 employed adults age 16 and older	NA	NA	` '	22.5 (2012)	
ercent of households that install a radon mitigation system when they receive a high radon test result	32% (2013-15)	39% (2014-16)	39% (2017)	NC	
ercent of riodscholds that install a lador mitigation system when they receive a high rador test result	NA	NA	17% (2017)	NA	
ercent of schools completing an environmental health walkinforgh	INA	INA	17/0 (2017)	IVA	
	NA		420/ (2015)	NI A	
ercent of inspections that find critical food safety violations	NA	NA	42% (2015)	NA NA	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters	NA 22.4 (2008)		42% (2015) 11.0 (2013)	NA NA	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING		NA	11.0 (2013)		
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent	22.4 (2008) NA	NA 19.2 (2011) NA	11.0 (2013) 49% (2013)	NA NC	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent	22.4 (2008)	NA 19.2 (2011)	11.0 (2013)	NA	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended	22.4 (2008) NA	NA 19.2 (2011) NA	11.0 (2013) 49% (2013)	NA NC	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent effectives ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE	22.4 (2008) NA 43% (2011) NA	NA 19.2 (2011) NA 50% (2015) NA	11.0 (2013) 49% (2013) 47% (2015) 87% (2016)	NC 27% NA	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent of ercent of high schools that teach key topics related to exemplary sexual health education in a required course ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE oronary heart disease death rate per 100,000 Vermonters	22.4 (2008) NA 43% (2011)	NA 19.2 (2011) NA 50% (2015) NA	11.0 (2013) 49% (2013) 47% (2015)	NA NC 27%	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE pronary heart disease death rate per 100,000 Vermonters	22.4 (2008) NA 43% (2011) NA	NA 19.2 (2011) NA 50% (2015) NA	11.0 (2013) 49% (2013) 47% (2015) 87% (2016)	NC 27% NA	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course  EART DISEASE & STROKE  Dronary heart disease death rate per 100,000 Vermonters  roke death rate per 100,000 Vermonters	22.4 (2008) NA 43% (2011) NA 102.8 (2008-10)	NA 19.2 (2011) NA 50% (2015) NA 105.6 (2012-14)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014)	NA  NC 27%  NA	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course  EART DISEASE & STROKE  Dronary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10)	NA 19.2 (2011) NA 50% (2015) NA 105.6 (2012-14) 30.3 (2012-14)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014)	NA  NC 27%  NA 98.8 36.5	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course  EART DISEASE & STROKE  Dronary heart disease death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IV & STD	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13) 73% (2011,13)	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 30.3 (2012-14) 27% (2013,15) 75% (2013,15)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015)	NA  NC 27%  NA  98.8 36.5 30% 76%	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE ercent of seast death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IV & STD  umber of new HIV diagnoses among all persons	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13) 73% (2011,13)	NA 19.2 (2011) NA 50% (2015) NA 105.6 (2012-14) 30.3 (2012-14) 27% (2013,15) 75% (2013,15)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015)	NA  NC 27%  NA 98.8 36.5 30% 76%	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE  Dronary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IV & STD  umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13) 73% (2011,13)  NA 6% (2011-12)	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 30.3 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2016)	NA  NC 27%  NA  98.8 36.5 30% 76%  NC 13%	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING  ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE  pronary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IV & STD  umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months ercent of adolescents in grades 9-12 ever tested for HIV	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13) 73% (2011,13)  NA 6% (2011-12) 12% (2009)	NA 19.2 (2011) NA 50% (2015) NA 105.6 (2012-14) 30.3 (2012-14) 27% (2013,15) 75% (2013,15)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2016) 10% (2015)	NA  NC  27%  NA  98.8  36.5  30%  76%  NC  13%  10%	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE  Dronary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IV & STD  umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months ercent of adolescents in grades 9-12 ever tested for HIV	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13) 73% (2011,13)  NA 6% (2011-12)	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 30.3 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2016)	NA  NC 27%  NA  98.8 36.5 30% 76%  NC 13%	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course  EART DISEASE & STROKE  Dronary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13) 73% (2011,13)  NA 6% (2011-12) 12% (2009)	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 30.3 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2016) 10% (2015)	NA  NC  27%  NA  98.8  36.5  30%  76%  NC  13%  10%	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE  Pronary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IV & STD  umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months ercent of adolescents in grades 9-12 ever tested for HIV ercent of persons linked to HIV care within 90 days of HIV diagnosis	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13) 73% (2011,13)  NA 6% (2011-12) 12% (2009) NA	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 30.3 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2016) 10% (2015) 90% (2015)	NA  NC  27%  NA  98.8  36.5  30%  76%  NC  13%  10%  NA	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING  ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course  EART DISEASE & STROKE  Pronary heart disease death rate per 100,000 Vermonters  Proke death rate per 100,000 Vermonters  Percent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IV & STD  Umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months ercent of adolescents in grades 9-12 ever tested for HIV ercent of persons linked to HIV care within 90 days of HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of gonorrhea per 100,000 females age 15-44	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13)  NA 6% (2011,13)  NA 6% (2011-12) 12% (2009) NA NA	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 30.3 (2012-14) 27% (2013,15)  NA 9% (2015-16) 13% (2015) NA NA	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015)	NA  NC  27%  NA  98.8  36.5  30%  76%  NC  13%  10%  NA  NA  263.4 (2015)	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course  EART DISEASE & STROKE  Dronary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IV & STD  Umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months ercent of adolescents in grades 9-12 ever tested for HIV ercent of persons linked to HIV care within 90 days of HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of gonorrhea per 100,000 females age 15-44 AMUNIZATION & INFECTIOUS DISEASE	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13)  NA 6% (2011,13)  12% (2009)  NA NA 22.7 (2006-10) 33.6 (2006-10)	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA NA 43.4 (2010-14) 37.8 (2010-14)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014)	NA  NC 27%  NA  98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING  ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE  Dronary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IV & STD  umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months ercent of adolescents in grades 9-12 ever tested for HIV ercent of persons linked to HIV care within 90 days of HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ate of gonorrhea per 100,000 females age 15-44 AMILY ALTION & INFECTIOUS DISEASE ercent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4)	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13)  NA 6% (2011,13)  12% (2009)  NA NA 22.7 (2006-10) 33.6 (2006-10)	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 30.3 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA NA 43.4 (2010-14)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014)	NA  NC  27%  NA  98.8  36.5  30%  76%  NC  13%  10%  NA  NA  263.4 (2015)	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE  Dronary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IV & STD  Lumber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months ercent of adolescents in grades 9-12 ever tested for HIV ercent of persons linked to HIV care within 90 days of HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of gonorrhea per 100,000 females age 15-44 atte of gonorrhea per 100,000 males age 15-44  MMUNIZATION & INFECTIOUS DISEASE	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13)  NA 6% (2011,13)  12% (2009)  NA NA 22.7 (2006-10) 33.6 (2006-10)	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA NA 43.4 (2010-14) 37.8 (2010-14)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014)	NA  NC 27%  NA  98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)	
ercent of inspections that find critical food safety violations eat stress related emergency department visits per 100,000 Vermonters  AMILY PLANNING ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE oronary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IV & STD  umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months ercent of adolescents in grades 9-12 ever tested for HIV ercent of persons linked to HIV care within 90 days of HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of gonorrhea per 100,000 females age 15-44 AMILY ALL ORDING SIFFECTIOUS DISEASE ercent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4) ercent of kindergarteners with 2 or more MMR doses	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13)  NA 6% (2011,13)  12% (2009)  NA NA 22.7 (2006-10) 33.6 (2006-10)	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 30.3 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA NA 43.4 (2010-14) 37.8 (2010-14)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)	
ercent of inspections that find critical food safety violations and stress related emergency department visits per 100,000 Vermonters  MILY PLANNING  Procent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE  Promonary heart disease death rate per 100,000 Vermonters Proceed eath rate per 100,000 Vermonters Procent of adults with hypertension Procent of adults with a cholesterol check in past 5 years  V & STD  Umber of new HIV diagnoses among all persons Procent of adults age 18-64 tested for HIV in past 12 months Procent of adolescents in grades 9-12 ever tested for HIV Procent of persons linked to HIV care within 90 days of HIV diagnosis Procent of persons with HIV viral suppression among those with HIV diagnosis Procent of gonorrhea per 100,000 females age 15-44 Procent of gonorrhea per 100,000 males age 15-44  PROMUNIZATION & INFECTIOUS DISEASE Procent of kindergarteners with 2 or more MMR doses Procent of adolescents age 13-15 who have completed the HPV vaccination series	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13)  NA 6% (2011,13)  12% (2009)  NA NA 22.7 (2006-10) 33.6 (2006-10)  NA NA	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% NA 10% NA 263.4 (2015) 307.6 (2015)	
recent of inspections that find critical food safety violations at stress related emergency department visits per 100,000 Vermonters  MILY PLANNING  From the of pregnancies that are intended erecent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse erecent of high schools that teach key topics related to exemplary sexual health education in a required course exercited to high schools that teach key topics related to exemplary sexual health education in a required course exercited to high schools that teach key topics related to exemplary sexual health education in a required course exercited to high schools that teach key topics related to exemplary sexual health education in a required course exercited to high schools that teach key topics related to exemplary sexual health education in a required course exercited for HISE and the exercited for adults with hypertension exercited for adults with hypertension exercited for adults with a cholesterol check in past 5 years  V & STD  Lordon of new HIV diagnoses among all persons exercited for HIV in past 12 months exercited for adults age 18-64 tested for HIV in past 12 months exercited for persons linked to HIV care within 90 days of HIV diagnosis exercited for persons with HIV viral suppression among those with HIV diagnosis exercited for persons with HIV viral suppression among those with HIV diagnosis exercited for persons with HIV viral suppression among those with HIV diagnosis exercited for persons with HIV viral suppression among those with HIV diagnosis exercited for persons with HIV viral suppression among those with HIV diagnosis exercited for persons with HIV viral suppression among those with HIV diagnosis exercited for persons with HIV viral suppression among those with HIV diagnosis exercited for persons with HIV viral suppression among those with HIV diagnosis exercited for persons with HIV viral suppression among those with HIV diagnosis exercited for HIV viral suppression among those with HIV	22.4 (2008)  NA  43% (2011)  NA  102.8 (2008-10)  35.1 (2008-10)  27% (2011,13)  NA  6% (2011-12)  12% (2009)  NA  NA  22.7 (2006-10)  33.6 (2006-10)  NA  22% (2013)	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 94% (2015)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015) 72% 95% NA 86%	
recent of inspections that find critical food safety violations at stress related emergency department visits per 100,000 Vermonters  MILY PLANNING  recent of pregnancies that are intended recent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent decourse recent of high schools that teach key topics related to exemplary sexual health education in a required course  FART DISEASE & STROKE  FOR ON ON VERMONTES  FOR ONE OF THE WARD ON THE W	22.4 (2008)  NA  43% (2011)  NA  102.8 (2008-10)  35.1 (2008-10)  27% (2011,13)  NA  6% (2011-12)  12% (2009)  NA  NA  22.7 (2006-10)  33.6 (2006-10)  35% (2010)  NA  22% (2013)  67% (2010)	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) 80% (2015)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 94% (2015) 94% (2015)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015) 72% 95% NA 86%	
ercent of inspections that find critical food safety violations at stress related emergency department visits per 100,000 Vermonters (MILY PLANNING)  ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course exact DISEASE & STROKE  romary heart disease death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  V & STD  umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months ercent of adolescents in grades 9-12 ever tested for HIV diagnosis ercent of persons linked to HIV care within 90 days of HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4) ercent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4) ercent of children age 13-15 who have completed the HPV vaccination series ercent of adolescents age 13-17 with at least 1 Tdap booster the of varicella (chicken pox) per 100,000 Vermonters age 17 or younger ercent of adults age 65 and older who receive annual flu shot ercent of adults age 65 and older who ever had pneumococcal vaccine	22.4 (2008)  NA  43% (2011)  NA  102.8 (2008-10)  35.1 (2008-10)  27% (2011,13)  NA  6% (2011-12)  12% (2009)  NA  NA  22.7 (2006-10)  33.6 (2006-10)  35% (2010)  NA  22% (2013)  67% (2010)  NA	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) NA	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 33% (2015) 96% (2015) 19.1 (2015)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015) 72% 95% NA 86% 197.0 (2013)	
recent of inspections that find critical food safety violations at stress related emergency department visits per 100,000 Vermonters  MILY PLANNING  Treent of pregnancies that are intended recent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse  Treent of high schools that teach key topics related to exemplary sexual health education in a required course in ISEASE & STROKE  Tronary heart disease death rate per 100,000 Vermonters  Treent of adults with hypertension  Treent of adults with a cholesterol check in past 5 years  V & STD  Treent of adults with a cholesterol check in past 12 months  Treent of adults age 18-64 tested for HIV in past 12 months  Treent of adolescents in grades 9-12 ever tested for HIV  Treent of persons linked to HIV care within 90 days of HIV diagnosis  Treent of persons with HIV viral suppression among those with HIV diagnosis  Treent of persons with HIV viral suppression among those with HIV diagnosis  Treent of persons with HIV viral suppression among those with HIV diagnosis  Treent of persons with HIV viral suppression among those with HIV diagnosis  Treent of persons with HIV viral suppression among those with HIV diagnosis  Treent of persons with HIV viral suppression among those with HIV diagnosis  Treent of persons with HIV viral suppression among those with HIV diagnosis  Treent of persons with HIV viral suppression among those with HIV diagnosis  Treent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4)  Treent of kindergarteners with 2 or more MMR doses  Treent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4)  Treent of kindergarteners with 2 or more MMR doses  Treent of adults age 65 and older who receive annual flushot  Treent of adults age 65 and older who receive annual flushot  Treent of adults age 65 and older who receive annual flushot  Treent of adults age 65 and older who receive annual flushot  Treent of adults age 65 and older who receive annual flushot  Tr	22.4 (2008)  NA  43% (2011)  NA  102.8 (2008-10)  35.1 (2008-10)  27% (2011,13)  NA  6% (2011-12)  12% (2009)  NA  NA  22.7 (2006-10)  33.6 (2006-10)  35% (2010)  NA  22% (2013)  67% (2010)  NA  66% (2011-12)	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) NA 62% (2015-16)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 33% (2015) 96% (2015) 19.1 (2015) 59% (2016)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015) 72% 95% NA 86% 197.0 (2013) 59% 72%	
ercent of inspections that find critical food safety violations at stress related emergency department visits per 100,000 Vermonters  IMILY PLANNING  From the of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse  From the of high schools that teach key topics related to exemplary sexual health education in a required course exert of high schools that teach key topics related to exemplary sexual health education in a required course exert DISEASE & STROKE  From the original properties of the properties of the properties of the properties of adults with hypertension ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  V & STD  Import of new HIV diagnoses among all persons  From the original properties of the propert	22.4 (2008)  NA  43% (2011)  NA  102.8 (2008-10)  35.1 (2008-10)  27% (2011,13)  NA  6% (2011-12)  12% (2009)  NA  NA  22.7 (2006-10)  33.6 (2006-10)  35% (2010)  NA  22% (2013)  67% (2010)  NA  66% (2011-12)  77% (2011-12)	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) NA 62% (2015-16) 72% (2015-16)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014)  76% (2015) 94% (2015) 33% (2015) 96% (2015) 19.1 (2015) 59% (2016) 77% (2016)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015) 72% 95% NA 86% 197.0 (2013) 59% 72%	
ercent of inspections that find critical food safety violations and stress related emergency department visits per 100,000 Vermonters  IMILY PLANNING  From to foregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE  Promary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IV & STD  umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months ercent of persons linked to HIV care within 90 days of HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4)  Percent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4)  Percent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4)  Percent of diagnoses age 13-17 with at least 1 Tdap booster  Recent o	22.4 (2008)  NA  43% (2011)  NA  102.8 (2008-10)  35.1 (2008-10)  27% (2011,13)  73% (2011,13)  NA  6% (2011-12)  12% (2009)  NA  22.7 (2006-10)  33.6 (2006-10)  35% (2010)  NA  22% (2013)  67% (2010)  NA  66% (2011-12)  77% (2011-12)  NA	NA 19.2 (2011)  NA 50% (2015) NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15) NA 9% (2015-16) 13% (2015) NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) NA 62% (2015-16) 72% (2015-16) NA	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 33% (2015) 96% (2015) 19.1 (2015) 59% (2016) 77% (2016) 100% (2017)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)  72% 95% NA 86% 197.0 (2013) 59% 72% 66% (2012)	
recent of inspections that find critical food safety violations at stress related emergency department visits per 100,000 Vermonters  IMILY PLANNING  Trent of pregnancies that are intended by the present of adolescents in grades 9-12 who used most or moderately effective contraception at most recent dercourse by the properties of high schools that teach key topics related to exemplary sexual health education in a required course by the promary heart disease death rate per 100,000 Vermonters by the promary heart disease death rate per 100,000 Vermonters by the deat	22.4 (2008)  NA  43% (2011)  NA  102.8 (2008-10)  35.1 (2008-10)  27% (2011,13)  73% (2011,13)  NA  6% (2011-12)  12% (2009)  NA  22.7 (2006-10)  33.6 (2006-10)  35% (2010)  NA  22% (2013)  67% (2010)  NA  66% (2011-12)  77% (2011-12)  NA	NA 19.2 (2011)  NA 50% (2015) NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15) NA 9% (2015-16) 13% (2015) NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) NA 62% (2015-16) 72% (2015-16) NA	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 33% (2015) 96% (2015) 19.1 (2015) 59% (2016) 77% (2016) 100% (2017)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)  72% 95% NA 86% 197.0 (2013) 59% 72% 66% (2012)	
ercent of inspections that find critical food safety violations and stress related emergency department visits per 100,000 Vermonters  VMILY PLANNING  Freent of pregnancies that are intended ercent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE  Formary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IN & STD  Umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months ercent of adolescents in grades 9-12 ever tested for HIV ercent of persons linked to HIV care within 90 days of HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with 10,000 females age 15-44  MMUNIZATION & INFECTIOUS DISEASE ercent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4) ercent of kindergarteners with 2 or more MMR doses ercent of adolescents age 13-15 with at least 1 Tdap booster of ercent of adolescents age 13-17 with at least 1 Tdap booster ercent of adolescents age 65 and older who receive annual flu shot ercent of adults age 65 and older who receive annual flu shot ercent of adults age 65 and older who receive annual flu shot ercent of adults age 65 and older who ever had pneumococcal vaccine ercent of adults age 65 and older who ever had pneumococcal vaccine ercent of adults age 65 and older who ever had pneumococcal vaccine ercent of drivers using car restraints	22.4 (2008)  NA  43% (2011)  NA  102.8 (2008-10)  35.1 (2008-10)  27% (2011,13)  NA  6% (2011-12)  12% (2009)  NA  NA  22.7 (2006-10)  35% (2010)  NA  22% (2013)  67% (2010)  NA  66% (2011-12)  77% (2011-12)  NA  NA  84% (2010)	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) NA 62% (2015-16) 72% (2015-16) NA NA	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 96% (2015) 19.1 (2015) 59% (2016) 77% (2016) 100% (2017) 1.005 (2016)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)  72% 95% NA 86% 197.0 (2013) 59% 72% 66% (2012) NA	
recent of inspections that find critical food safety violations and stress related emergency department visits per 100,000 Vermonters  MILY PLANNING  The control of pregnancies that are intended erecent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse erecent of high schools that teach key topics related to exemplary sexual health education in a required course each of high schools that teach key topics related to exemplary sexual health education in a required course each of high schools that teach key topics related to exemplary sexual health education in a required course each of high schools that teach key topics related to exemplary sexual health education in a required course each of high schools that teach key topics related to exemplary sexual health education in a required course each execution of high schools that teach key topics related to exemplary sexual health education in a required course each execution of high schools that teach key topics related to exemplary sexual health education in a required course each execution of high schools that teach key topics related to exemplary sexual health education in a required course each execution at the execution of adults with a cholesterol check in past 5 years  I was stored of adults age 18-64 tested for HIV in past 12 months executed for HIV diagnosis executed persons with HIV viral suppression among those with HIV diagnosis executed persons with HIV viral suppression among those with HIV diagnosis executed for provide age 15-44  Intendict of persons with HIV viral suppression among those with HIV diagnosis executed for foliopen age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4)  Intendict of persons with execution and prevention series executed for HIV executed and trien to incentine active 18 case to make a prevention of the fol	22.4 (2008)  NA  43% (2011)  NA  102.8 (2008-10)  35.1 (2008-10)  27% (2011,13)  NA  6% (2011-12)  12% (2009)  NA  NA  22.7 (2006-10)  33.6 (2006-10)  35% (2010)  NA  22% (2013)  67% (2010)  NA  66% (2011-12)  77% (2011-12)  NA  NA  84% (2010)  926.1 (2008-10)	NA 19.2 (2011)  NA 50% (2015)  NA 105.6 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) NA 62% (2015-16) 72% (2015-16) 72% (2015-16) NA NA	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10 (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 33% (2015) 96% (2015) 19.1 (2015) 59% (2016) 77% (2016) 100% (2017) 1.005 (2016) 85% (2015)	NA  NC 27% NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)  72% 95% NA 86% 197.0 (2013) 59% 72% 66% (2012) NA NA 736.1 (2013) 5474.6	
recent of inspections that find critical food safety violations that stress related emergency department visits per 100,000 Vermonters  MILY PLANNING  recent of pregnancies that are intended recent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse recent of high schools that teach key topics related to exemplary sexual health education in a required course  ART DISEASE & STROKE  Bronary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters recent of adults with hypertension recent of adults with a cholesterol check in past 5 years  V & STD  Imber of new HIV diagnoses among all persons recent of adults age 18-64 tested for HIV in past 12 months recent of adults age 18-64 tested for HIV in past 12 months recent of persons linked to HIV care within 90 days of HIV diagnosis recent of persons with HIV viral suppression among those with HIV diagnosis recent of persons with HIV viral suppression among those with HIV diagnosis recent of persons with HIV viral suppression among those with HIV diagnosis recent of persons with HIV viral suppression among those with HIV diagnosis recent of persons with HIV viral suppression among those with HIV diagnosis recent of persons with HIV viral suppression among those with HIV diagnosis recent of persons with HIV viral suppression among those with HIV diagnosis recent of persons with HIV viral suppression among those with HIV diagnosis recent of of persons with HIV viral suppression among those with HIV diagnosis recent of of persons with HIV viral suppression among those with HIV diagnosis recent of of persons with HIV viral suppression among those with HIV diagnosis recent of of of persons with HIV viral suppression among those with HIV diagnosis recent of adolescents age 13-15 who have completed the HPV vaccination series recent of adolescents age 13-15 who have completed the HPV vaccination series recent of adolescents age 13-17 with at least 1 Tdap booster te of varicella (chicken pox) p	22.4 (2008)  NA  43% (2011)  NA  102.8 (2008-10)  35.1 (2008-10)  27% (2011,13)  NA  6% (2011-12)  12% (2009)  NA  NA  22.7 (2006-10)  33.6 (2006-10)  35% (2010)  NA  22% (2013)  67% (2010)  NA  66% (2011-12)  77% (2011-12)  NA  NA  84% (2010)  926.1 (2008-10)	NA 19.2 (2011)  NA 50% (2015) NA 105.6 (2012-14) 27% (2013,15) 75% (2013,15) NA 9% (2015-16) 13% (2015- NA NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) NA 62% (2015-16) 72% (2015-16) 72% (2015-16) NA NA 87% (2014) 962.9 (2009-11)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 33% (2015) 96% (2015) 19.1 (2015) 59% (2016) 77% (2016) 100% (2017) 1.005 (2016) 85% (2015) 860.1 (2011)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015) 72% 95% NA 86% 197.0 (2013) 59% 72% 66% (2012) NA NA 736.1 (2013)	
Arcent of inspections that find critical food safety violations at stress related emergency department visits per 100,000 Vermonters (MMILY PLANNING)  Arcent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse  Foreign of high schools that teach key topics related to exemplary sexual health education in a required course tercourse  FORTH DISEASE & STROKE  FORTH PROPERTY HEADTH OF THE	22.4 (2008)  NA  43% (2011)  NA  102.8 (2008-10)  35.1 (2008-10)  27% (2011,13)  73% (2011,13)  NA  6% (2011-12)  12% (2009)  NA  22.7 (2006-10)  33.6 (2006-10)  35% (2010)  NA  22% (2013)  67% (2010)  NA  66% (2011-12)  77% (2011-12)  NA  NA  84% (2010)  926.1 (2008-10)  NA  165.3 (2008-10)	NA 19.2 (2011)  NA 50% (2015) NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15) 75% (2015-16) 13% (2015- NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) 80% (2015) 80% (2015-16) 72% (2015-16) NA NA 87% (2014) 962.9 (2009-11) 6369.2 (2009-11) 57.1 (2012-14)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 33% (2015) 96% (2015) 19.1 (2015) 59% (2016) 77% (2016) 100% (2017) 1.005 (2016) 85% (2015) 860.1 (2011) 5628.4 (2011) 118.7 (2014)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)  72% 95% NA 86% 197.0 (2013) 599% 66% (2012) NA  NA 736.1 (2013) 5474.6 (2012) 58.8	
ercent of inspections that find critical food safety violations and stress related emergency department visits per 100,000 Vermonters  MILY PLANNING  recent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE  pronary heart disease death rate per 100,000 Vermonters roke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  IV & STD  umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months ercent of adolescents in grades 9-12 ever tested for HIV ercent of persons linked to HIV care within 90 days of HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV wiral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4)  **HIMUNIZATION & INFECTIOUS DISEASE**  ercent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4)  **Ercent of adolescents age 13-17 with at least 1 Tdap booster ercent of adolescents age 13-17 with at least 1 Tdap booster ercent of adolescents age 13-17 with at least 1 Tdap booster ercent of adults age 65 and older who ever had pneumococcal vaccine ercent of adults age 65 and older who ever had pneumococcal vaccine ercent of adults age 65 and older who ever had pneumococcal vaccine ercent of adults age 65 and older who ever had pneumococcal vaccine ercent of adults age 65 and older who ever had pneumococcal vaccine ercent of adults age 65 and older who ever had pneumococcal vaccine ercent of adults age 65 and older who e	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13) 73% (2011,13)  NA 6% (2011-12) 12% (2009) NA NA 22.7 (2006-10) 33.6 (2006-10)  35% (2010) NA 22% (2013) 67% (2010) NA 66% (2011-12) 77% (2011-12) NA NA  84% (2010) 926.1 (2008-10) NA 165.3 (2008-10) 238.4 (2008-10)	NA 19.2 (2011)  NA 50% (2015) NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15) 75% (2015-16) 13% (2015- NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) 80% (2015) 80% (2015- 16) 72% (2015-16) 72% (2015-16) NA 87% (2014) 962.9 (2009-11) 57.1 (2012-14) 247.5 (2009-11)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 33% (2015) 96% (2015) 19.1 (2015) 59% (2016) 77% (2016) 100% (2017) 1.005 (2016) 85% (2015) 860.1 (2011) 5628.4 (2011) 118.7 (2014) 173.1 (2011)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)  72% 95% NA 86% 197.0 (2013) 599% 66% (2012) NA  NA 736.1 (2013) 5474.6 (2012) 58.8 160.5 (2013)	4
Arcent of inspections that find critical food safety violations are stress related emergency department visits per 100,000 Vermonters  MMILY PLANNING  Arcent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse  Frecent of high schools that teach key topics related to exemplary sexual health education in a required course tercourse  Frecent of high schools that teach key topics related to exemplary sexual health education in a required course teach that the per 100,000 Vermonters roke death rate per 100,000 Vermonters  Frecent of adults with hypertension ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  V & STD  Wimber of new HIV diagnoses among all persons  Frecent of adults age 18-64 tested for HIV in past 12 months  Frecent of adults age 18-64 tested for HIV in past 12 months  Frecent of persons linked to HIV care within 90 days of HIV diagnosis  Frecent of persons with HIV viral suppression among those with HIV diagnosis  Frecent of persons with HIV viral suppression among those with HIV diagnosis  Frecent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4)  Frecent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4)  Frecent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4)  Frecent of adolescents age 13-15 who have completed the HPV vaccination series  Frecent of adults age 65 and older who receive annual flu shot  Frecent of adults age 65 and older who receive annual flu shot  Frecent of adults age 65 and older who receive annual flu shot  Frecent of orders using car restraints  Frecent of adults age 65 and older who receive annual flu shot  Frecent of orders using car restraints  Frecent of divers using car restraints  Frecent of children ratio for Clostridium difficile infections (C. diff)  Frecent of divers using car restraints  Frecent of divers using car restraints  Frecent of divers using c	22.4 (2008)  NA  43% (2011)  NA  102.8 (2008-10)  35.1 (2008-10)  27% (2011,13)  73% (2011,13)  NA  6% (2011-12)  12% (2009)  NA  22.7 (2006-10)  33.6 (2006-10)  35% (2010)  NA  22% (2013)  67% (2010)  NA  66% (2011-12)  77% (2011-12)  NA  NA  84% (2010)  926.1 (2008-10)  NA  165.3 (2008-10)	NA 19.2 (2011)  NA 50% (2015) NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15) 75% (2015-16) 13% (2015- NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) 80% (2015) 80% (2015-16) 72% (2015-16) NA NA 87% (2014) 962.9 (2009-11) 6369.2 (2009-11) 57.1 (2012-14)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 33% (2015) 96% (2015) 19.1 (2015) 59% (2016) 77% (2016) 100% (2017) 1.005 (2016) 85% (2015) 860.1 (2011) 5628.4 (2011) 118.7 (2014)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)  72% 95% NA 86% 197.0 (2013) 599% 66% (2012) NA  NA 736.1 (2013) 5474.6 (2012) 58.8	
where the finispections that find critical food safety violations are stress related emergency department visits per 100,000 Vermonters  MINITY PLANNING  Incent of pregnancies that are intended executed advised in grades 9-12 who used most or moderately effective contraception at most recent derectourse executed in grades 9-12 who used most or moderately effective contraception at most recent derectourse executed in grades 9-12 who used most or moderately effective contraception at most recent derectourse executed for grades and the grades of	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13) 73% (2011,13)  NA 6% (2011-12) 12% (2009) NA NA 22.7 (2006-10) 33.6 (2006-10)  35% (2010) NA 22% (2013) 67% (2010) NA 66% (2011-12) 77% (2011-12) NA NA  84% (2010) 926.1 (2008-10) NA 165.3 (2008-10) 238.4 (2008-10)	NA 19.2 (2011)  NA 50% (2015) NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15) 75% (2015-16) 13% (2015- NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) 80% (2015) 80% (2015- 16) 72% (2015-16) 72% (2015-16) NA 87% (2014) 962.9 (2009-11) 57.1 (2012-14) 247.5 (2009-11)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 33% (2015) 96% (2015) 19.1 (2015) 59% (2016) 77% (2016) 100% (2017) 1.005 (2016) 85% (2015) 860.1 (2011) 5628.4 (2011) 118.7 (2014) 173.1 (2011)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)  72% 95% NA 86% 197.0 (2013) 59% 72% 66% (2012) NA  NA 736.1 (2013) 5474.6 (2012) 58.8 160.5 (2013) 10.3	
Arcent of inspections that find critical food safety violations at stress related emergency department visits per 100,000 Vermonters  MINITY PLANNING  Treent of pregnancies that are intended erecent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse  Treent of high schools that teach key topics related to exemplary sexual health education in a required course exercity of the properties	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13) 73% (2011,13)  NA 6% (2011-12) 12% (2009) NA NA 22.7 (2006-10) 33.6 (2006-10)  35% (2010) NA 22% (2013) 67% (2010) NA 66% (2011-12) 77% (2011-12) NA NA  84% (2010) 926.1 (2008-10) 238.4 (2008-10) 9.5 (2009-11) NA	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) 80% (2015) 80% (2015-16) 72% (2015-16) NA NA 87% (2014) 962.9 (2009-11) 57.1 (2012-14) 247.5 (2009-11) 9.9 (2012-14)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 33% (2015) 96% (2015) 19.1 (2015) 59% (2016) 77% (2016) 100% (2017) 1.005 (2016) 85% (2015) 860.1 (2011) 5628.4 (2011) 118.7 (2014) 173.1 (2011) 9.5 (2014)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)  72% 95% NA 86% 197.0 (2013) 59% 72% 66% (2012) NA  NA 736.1 (2013) 58.8 160.5 (2013) 10.3	
Arcent of inspections that find critical food safety violations at stress related emergency department visits per 100,000 Vermonters  MINITY PLANNING  Procent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse ercent of high schools that teach key topics related to exemplary sexual health education in a required course EART DISEASE & STROKE  Toronary heart disease death rate per 100,000 Vermonters oroke death rate per 100,000 Vermonters ercent of adults with hypertension ercent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  V & STD  Umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months ercent of adults age 18-64 tested for HIV in past 12 months ercent of adults age 18-64 tested for HIV in past 12 months ercent of persons linked to HIV care within 90 days of HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of persons with HIV viral suppression among those with HIV diagnosis ercent of children age 19-35 months receiving recommended vaccines (4:3:1:4:3:1:4) ercent of kindergarteners with 2 or more MMR doses  Percent of kindergarteners with 2 or more MMR doses  Percent of adolescents age 13-15 who have completed the HPV vaccination series ercent of adults age 65 and older who ever had pneumococcal vaccine ercent of adults age 65 and older who ever had pneumococcal vaccine ercent of adults age 65 and older who ever had pneumococcal vaccine ercent of incument active 10 case contacts with newly-diagnose LTD wino Started and then completed adults and for Clostridium difficile infections (C. diff)  DURY & VIOLENCE PREVENTION  Percent of divers using car restraints of the percent of adults age 65 and older who ever had pneumococcal vaccine ercent of didvers using car restraints of the percent of the presence of the	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13) 73% (2011,13)  NA 6% (2011-12) 12% (2009) NA NA 22.7 (2006-10) 33.6 (2006-10)  35% (2010) NA 22% (2013) 67% (2010) NA 66% (2011-12) 77% (2011-12) NA NA  84% (2010) 926.1 (2008-10) 238.4 (2008-10) 9.5 (2009-11)  NA NA NA	NA 19.2 (2011)  NA 50% (2015) NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15) 75% (2015-16) 13% (2015- NA NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015- NA 62% (2015- NA 62% (2015- NA 62% (2015- NA NA 87% (2014) 962.9 (2009-11) 57.1 (2012-14) 247.5 (2009-11) 9.9 (2012-14)  NA NA	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 33% (2015) 96% (2015) 19.1 (2015) 59% (2016) 77% (2016) 100% (2017) 1.005 (2016) 85% (2015) 860.1 (2011) 5628.4 (2011) 118.7 (2014) 173.1 (2011) 9.5 (2014) 0.44 (2011-13) 87% (2013)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)  72% 95% NA 86% 197.0 (2013) 599% 66% (2012) NA  NA 736.1 (2013) 5474.6 (2012) 58.8 160.5 (2013) 10.3 0.87 (2013) 92%	4
Arrent of inspections that find critical food safety violations at stress related emergency department visits per 100,000 Vermonters  MMILY PLANNING  Percent of pregnancies that are intended ercent of adolescents in grades 9-12 who used most or moderately effective contraception at most recent tercourse  Percent of injsh schools that teach key topics related to exemplary sexual health education in a required course tercourse  Percent of injsh schools that teach key topics related to exemplary sexual health education in a required course tercent of injsh schools that teach key topics related to exemplary sexual health education in a required course teach recent of sexual sease death rate per 100,000 Vermonters or the death rate per 100,000 Vermonters  Percent of adults with hypertension ercent of adults with a cholesterol check in past 5 years  V & STD  Umber of new HIV diagnoses among all persons ercent of adults age 18-64 tested for HIV in past 12 months  Percent of persons linked to HIV care within 90 days of HIV diagnosis  Percent of persons with HIV viral suppression among those with HIV diagnosis ercent of genorrhea per 100,000 females age 15-44  MINUNIZATION & INFECTIOUS DISEASE  Percent of Kindergarteners with 2 or more MMR doses  Percent of Adolescents age 13-15 who have completed the HPV vaccination series  Percent of Adolescents age 13-15 who have completed the HPV vaccination series  Percent of adolescents age 13-17 with at least 1 Tdap booster  Percent of adolescents age 13-17 with at least 1 Tdap booster  Percent of adolescents age 13-17 with at least 1 Tdap booster  Percent of adolescents age 13-15 who have completed the HPV vaccination series  Percent of adolescents age 13-15 with at least 1 Tdap booster  Percent of adolescents age 13-15 with at least 1 Tdap booster  Percent of adolescents age 13-15 with a least 1 Tdap booster  Percent of adolescents age 13-15 with a least 1 Tdap booster  Percent of adolescents age 13-15 with a least 1 Tdap booster  Percent of adolescents age 13-15 with a least 1	22.4 (2008)  NA 43% (2011)  NA  102.8 (2008-10) 35.1 (2008-10) 27% (2011,13) 73% (2011,13)  NA 6% (2011-12) 12% (2009) NA NA 22.7 (2006-10) 33.6 (2006-10)  35% (2010) NA 22% (2013) 67% (2010) NA 66% (2011-12) 77% (2011-12) NA NA  84% (2010) 926.1 (2008-10) 238.4 (2008-10) 9.5 (2009-11) NA	NA 19.2 (2011)  NA 50% (2015)  NA  105.6 (2012-14) 27% (2013,15) 75% (2013,15)  NA 9% (2015-16) 13% (2015) NA 43.4 (2010-14) 37.8 (2010-14) 64% (2015) NA 26% (2015) 80% (2015) 80% (2015-16) 72% (2015-16) NA NA 87% (2014) 962.9 (2009-11) 57.1 (2012-14) 247.5 (2009-11) 9.9 (2012-14)	11.0 (2013) 49% (2013) 47% (2015) 87% (2016) 105.4 (2014) 27.7 (2014) 25% (2015) 76% (2015) 10% (2016) 10% (2015) 90% (2015) 94% (2015) 30.2 (2014) 41.6 (2014) 76% (2015) 94% (2015) 33% (2015) 96% (2015) 19.1 (2015) 59% (2016) 77% (2016) 100% (2017) 1.005 (2016) 85% (2015) 860.1 (2011) 5628.4 (2011) 118.7 (2014) 173.1 (2011) 9.5 (2014)	NA  NC 27%  NA 98.8 36.5 30% 76%  NC 13% 10% NA NA 263.4 (2015) 307.6 (2015)  72% 95% NA 86% 197.0 (2013) 599% 66% (2012) NA  NA 736.1 (2013) 5474.6 (2012) 58.8 160.5 (2013) 10.3 0.87 (2013) 92%	

http://healthvermont.gov/about/reports/healthy-vermonters-plans-reports

VERMONT DEPARTMENT OF HEALTH

## Healthy Vermonters 2020 - Windham County Quick Reference

Healthy Vermonters 2020 Indicator	County Baseline	County Current	VT Current	U.S. Current	Targ
MATERNAL & INFANT HEALTH continued from page 1					
Percent women delivering a live birth who discussed preconception health prior to pregnancy	NA	NA	21% (2013)	23%	
Percent of women delivering a live birth who had a healthy weight prior to pregnancy	NA	NA NA	46% (2013)		
Percent of women who had a dental visit during pregnancy	NA	NA NA	59% (2013)		
Percent of infants breastfed exclusively for six months Percent of children who live in households where someone smokes	NA NA	NA NA	28% (2014) 19% (2016)	22% 16%	
MENTAL HEALTH	INA	IVA	1970 (2010)	10/0	, 1
Rate of suicide per 100,000 Vermonters	15.4 (2008-10)	20.5 (2012-14)	17.2 (2014)	13.0	) 1
Percent of adolescents in grades 9-12 who made a suicide plan	12% (2011)	14% (2015)	12% (2015)	15%	, )
Percent of Medicaid beneficiaries age 10-17 screened for depression	NA	26% (2015-16)	18% (2015)	NA	
Percent of Medicaid beneficiaries age 18 and older screened for depression	NA	5% (2015-16)	5% (2015)	NA	
Percent of Vermont adults with any mental health conditions receiving treatment	NA	NA	56% (2016)	45%	,
NUTRITION & WEIGHT STATUS					
Percent of adults age 20 and older who are obese	23% (2011-12)	25% (2015-16)	28% (2016)	30%	
Percent of children age 2-5 (in WIC) who are obese	12% (2011)	14% (2014)		14% (2010)	
Percent of adolescents in grades 9-12 who are obese	11% (2011)	12% (2015)	12% (2015)	14%	
Percent of households with food insecurity	NA	NA		14% (2014)	
Percent of adults eating fruit 2 or more times daily	34% (2011,13)	34% (2013,15)	32% (2015)		
Percent of adolescents in grades 9-12 eating fruit 2 or more times daily	31% (2011)	31% (2015)	34% (2015)		
Percent of adults eating the daily vegetables 3 or more times daily	19% (2011,13)	21% (2013,15)	20% (2015)	17%	
Percent of adolescents in grades 9-12 eating vegetables 3 or more time daily	14% (2011)	14% (2015)	18% (2015)	15%	5 2
Percent of adults consuming less than 1 sugar sweetened beverage per day	NA	NA	83% (2013)	NA	. 8
Percent of adolescents in grades 9-12 consuming less than 1 sugar sweetened beverage per day	69% (2011)	76% (2015)	77% (2015)	NA	. 8
DIDER ADULTS  Percent of adults are SE and older who visited a dector in the last year for routine care.	940/ /2044 421	959/ (2045-46)	000/ /2040	000/ /2045	
Percent of adults age 65 and older who visited a doctor in the last year for routine care	84% (2011-12) 56% (2008 10)	85% (2015-16) 45% (2014-16)		88% (2015)	
ercent of males age 65 and older who are up to date on a core set of clinical preventive services	56% (2008,10)	45% (2014,16)		40% (2015)	
Percent of females age 65 and older who are up to date on a core set of clinical preventive services	46% (2008,10)	38% (2012,14) NA		35% (2015)	
ercent of adults age 65 and older with prescription drug coverage  ORAL HEALTH	NA	NA	84% (2012)	85%	•
ercent of children age 6-9 with dental decay	NA	NA	31% (2017)	NC	
ercent of adults age 45-64 with tooth extraction	52% (2006,08,10)	50% (2014,16)	49% (2016)	54%	
Percent of children in grades K-6 using dental system yearly	NA	NA	77% (2016)		
Percent of children in grades 7-12 using dental system yearly	NA NA	NA NA	61% (2016)	NA	
ercent of adults using dental system yearly	70% (2006,08,10)	73% (2014,16)	71% (2014)	65%	
ercent of addits using deficit system yearry ercent of population served by community public water systems that have optimally fluoridated water	NA	7578 (2014,10) NA	56% (2015)	75%	
PHYSICAL ACTIVITY	IVA	IVA	30% (2013)	75/0	,
ercent of adults with no leisure time aerobic physical activity	15% (2011-12)	18% (2014-15)	18% (2016)	24%	,
Percent of adults meeting aerobic physical activity guidelines	60% (2011,13)	60% (2013,15)	59% (2015)	51%	,
ercent of adolescents in grades 9-12 meeting physical activity guidelines	24% (2011)	21% (2015)	23% (2015)	27%	,
ercent of adolescents in grades 9-12 who did not participate in physical activity for at least 60 minutes on any	15% (2011)	15% (2015)	14% (2015)	14%	,
day during the past 7 days PUBLIC HEALTH PREPAREDNESS	1370 (2011)	1370 (2013)	1470 (2013)	1470	,
Time necessary to issue official information regarding a public health emergency	NA	NA	60 min	NΛ	60
Time necessary to activate personnel for a public health emergency	NA NA	NA NA	28 min (2014)		
Percent of after-action reports and improvement plans that are drafted and submitted in 60 days or less					
ollowing an emergency & exercises	NA	NA	95% (2016)	NC	
roportion of crisis and emergency risk messages intended to protect the public's health that demonstrate the	NA	NA	75% (2016)	NA	. 1
ise of best practices IESPIRATORY DISEASES					
Asthma hospitalization rate per 10,000 children age 4 or younger	17.0 (2008-10)	20.5 (2009-11)	10.6 (2011)	40.6 (2009)	) :
sthma hospitalization rate per 10,000 Vermonters age 5-64	3.6 (2008-10)	3.3 (2009-11)		10.5 (2010)	
sthma hospitalization rate per 10,000 adults age 65 and older	4.9 (2008-10)	6.3 (2009-11)		25.5 (2010)	
ercent of adult non-smokers exposed to secondhand smoke	52% (2008,10)	48% (2014,16)	44% (2016)		
Percent of adults with asthma who have a written asthma management plan from a doctor or health care					
rofessional	20% (2008-10)	17% (20012-14)	33% (2014)	31% (2013)	)
ercent of children age 17 or younger with asthma who have a written asthma management plan from a doctor	, NA	NA	48% (2010)	49%	
or health care professional					
ercent of adults with asthma who are advised to change things in their environment	48% (2008-10)	59% (2012-14)	38% (2014)		
ercent of children age 17 or younger with asthma who are advised to change things in their environment  CHOOL AGE HEALTH	NA	NA	33% (2010)	36%	5
ercent of kindergarteners ready for school in all five domains of healthy development	NA	NA	82% (2015-16)	NA	
ercent of students age 10-17 who have had a wellness exam in past 12 months	NA NA	NA NA	65% (2015-16)		
ercent of schools that provide referrals to community based medical providers for students identified with					
hronic conditions or at risk for activity, diet and weight-related chronic conditions (all schools grades 6-12)	NA	NA	82% (2016)	55%	•
OCIAL DETERMINANTS OF HEALTH					
ercent of adults with disabilities always or usually getting social and emotional support	77% (2008-10)	72% (2012,14)	74% (2014)	72% (2015)	)
ercent of Vermonters living below the poverty level	11% (2006-10)	13% (2011-15)	12% (2011-15)	15%	ò
ercent of Vermonters age 17 or younger living below the poverty level	13% (2006-11)	17% (2011-15)	15% (2011-15)	22%	ò
ercent of households that spend 30% or more of their income on housing	38% (2006-11)	40% (2011-15)	37% (2011-15)	35%	)
ercent of students awarded a high school diploma 4 years after starting 9th grade	NA	NA	88% (2013-14)	82%	)
ercent of eligible registered voters who participated in the last election	73% (2008)	69% (2016)	68% (2016)	60%	5
JBSTANCE ABUSE					
ercent of persons age 12 and older who need and do not receive alcohol treatment	NA	NA	7% (2013-14)		
ercent of persons age 12 and older who need and do not receive illicit drug use treatment	NA	NA	3% (2013-14)	2%	
ercent of adolescents in grades 9-12 who used marijuana in the past 30 days	29% (2011)	26% (2015)	22% (2015)	22%	
ercent of adolescents in grades 9-12 binge drinking in the past 30 days	22% (2011)	16% (2015)	16% (2015)	18%	
ercent of adults age 18-24 binge drinking in the last 30 days	NA	18% (2014-2016)	32% (2016)	25% (2015)	)
ercent of adults age 65 and older who drink at a level of risk	26% (2011-12)	29% (2015-16)	23% (2016)	18% (2015)	)
		2021 15 2 1	.=		
	18% (2011-12)	23% (2014-15)	17% (2015)	17%	
ercent of adults who smoke cigarettes			110/ /2014\	NC	:
ercent of adults who smoke cigarettes ercent of adults using smokeless or other tobacco products	NA	12% (2012-14)	11% (2014)		
ercent of adults who smoke cigarettes ercent of adults using smokeless or other tobacco products ercent of adolescents in grades 9-12 who smoke cigarettes		11% (2015)	11% (2015)	11%	
ercent of adults who smoke cigarettes ercent of adults using smokeless or other tobacco products ercent of adolescents in grades 9-12 who smoke cigarettes ercent of adolescents in grades 9-12 who use e-cigarettes	NA	11% (2015) 13% (2015)	11% (2015) 15% (2015)		
OBACCO USE ercent of adults who smoke cigarettes ercent of adults using smokeless or other tobacco products ercent of adolescents in grades 9-12 who smoke cigarettes ercent of adolescents in grades 9-12 who use e-cigarettes ercent of adult smokers who attempted to quit smoking in the past year	NA 18% (2011)	11% (2015)	11% (2015)	11%	, )

Notes:

1/31/2018

**VERMONT** 

DEPARTMENT OF HEALTH