
2011 Vermont Youth Risk Behavior Survey Report

Report for Washington South Supervisory Union

Every other year, since 1993, the Department of Health Division of Alcohol and Drug Abuse Programs and the Department of Education Student Health and Learning team sponsor the Vermont Youth Risk Behavior Survey (YRBS). The YRBS measures the prevalence of behaviors that contribute to the leading causes of death, disease, and injury among youth.

Participation by schools and individual students is voluntary. To protect student privacy, the questionnaire and survey process are anonymous. It is impossible to identify an individual student's responses.

Survey Changes in 2011

Before 2011, students in grades 8-12 took the Vermont Youth Risk Behavior Survey. In 2011, we conducted two surveys: a **high school survey** of students in grades 9-12, and a **middle school survey** of students in grades 6-8.

All results in the high school survey report are for grades 9-12 only. For the questions on the 2011 high school survey, we recalculated results from previous years for grades 9-12 for comparative purposes. Those results are at the end of this report.

About this Report

The results appear in data tables by gender and grade unless otherwise noted. All results are percentages of students who responded affirmatively. Percentages were suppressed with an asterisk (*) if five or fewer students reported affirmatively, either overall or by grade or gender.

Thank You

We thank the principals and superintendents who chose to participate, the teachers and school staff who administered or in other ways supported the survey, and the students who took the time and effort to share a piece of their lives with us. We also thank the Centers for Disease Control and Prevention, Division of Adolescent and School Health for sponsoring the survey through a cooperative agreement with the Vermont Department of Education, and the CDC and Westat for technical assistance.

Table of Contents

BACKGROUND INFORMATION	
How to Use the YRBS	3
A Word of Caution	4
About the Surveys	5
HIGH SCHOOL SURVEY RESULTS	15
Sample Demographics	16
Personal Safety	17
Alcohol, Tobacco, and Other Drug Use	21
Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use.....	28
Sexual Behavior and Orientation	30
Body Image	33
Nutrition.....	35
Physical Activity.....	36
Youth Assets	37
MIDDLE SCHOOL SURVEY RESULTS.....	39
Sample Demographics	40
Personal Safety	41
Alcohol, Tobacco, and Other Drug Use	45
Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use.....	49
Body Image	51
Nutrition.....	52
Physical Activity.....	53
Youth Assets	55
RECALCULATED HIGH SCHOOL RESULTS	57

How to Use the YRBS

The YRBS can detect changes in risk behaviors over time and identify differences among ages, grades, and genders. With these data, we can focus prevention efforts and determine whether school policies and community programs are having the intended effect on student behaviors.

Think of the YRBS as a tool for starting discussions, for educating the community, for planning and evaluating programs, and for comparing Vermont students with other students nationwide.

- **Start the Conversation:** Use the YRBS to begin a conversation with teens about the personal choices they make or about the health of their community. Ask them if the results accurately reflect what they see happening around them. How do they explain the results? From their perspective, what is or is not working? How would they promote healthy behaviors?
- **Increase Awareness:** The YRBS provides an opportunity to break through “denial” and make community members aware of the risks that their young people face. It can also dispel myths and correct misinformation about the “average teenager.” The YRBS can accentuate the positive and celebrate the fact that many students are abstaining from behaviors that endanger their health and their ability to succeed.
- **Plan and Evaluate Programs:** The YRBS can serve as the basis of a community needs assessment. It can help identify strengths and weaknesses in communities, and can inform strategies to address those weaknesses.

Remember to Look at the Positive Side: In most cases, the majority of adolescents are NOT engaging in risky behaviors. Although most of the charts examine the prevalence of risk behaviors, please do not forget about the percentage of adolescents who are NOT engaging in these behaviors.

A Word of Caution

The YRBS represents the most complete and most recent information available about risk behaviors among Vermont students. However, the YRBS has some limitations that you should keep in mind when interpreting the results.

- **Data Quality:** We take numerous precautions to ensure the reliability and validity of the results. The Centers for Disease Control and Prevention (CDC) carefully designed and thoroughly tested the questionnaire. The anonymous survey encourages students to be honest and forthright. We run over 100 consistency checks on the data to exclude careless, invalid, or logically inconsistent answers. These precautions can reduce some sources of error, but not all.
- **Comparing Supervisory Unions/School Districts to Each Other and to the State:** It is natural to want to know how individual supervisory unions or school districts compare to the state overall or to each other. We urge caution in making such comparisons because the statewide results are weighted and the local results are not. As a result, it is possible that apparent differences, especially small differences, are due to demographic characteristics rather than to true differences in prevalence. Furthermore, small differences may not be statistically different and may simply be a function of normal sampling error.
- **What, not Why:** The YRBS can indicate what students are doing. It can also suggest which groups of students are more likely to engage in these behaviors. However, the survey does not answer the most important question: Why are they doing it?

Personal Safety

Feeling safe – whether at school, in a car, or in a relationship – fosters positive adolescent development. The high school and middle school surveys included questions on fighting, bullying, seat belt use, safety at school, helmet use, and suicidality. The high school survey included additional questions on driving behaviors, dating violence, and self harm.

- Physical fighting is a marker for problem behaviors¹ and is associated with serious injury.^{2,3} Abuse by an intimate partner and forced sex are associated with negative psychosocial outcomes, poor mental health outcomes, and other risk behaviors among both males and females.^{4,5,6}
- Both being a bully and being victimized by bullies have been increasingly recognized as health problems for children because of their association with a range of problems, including poor psychological adjustment,^{7,8} poor academic achievement,⁸ and violent behavior.⁹
- Motor vehicle crash injuries are the leading cause of death among youth ages 15 to 19 in the U.S.¹⁰ Proper use of safety belts reduces the risk of fatal injury to front seat passengers by 45% and risk of moderate to critical injury by 50%.¹¹
- Bicycle helmets are 85% to 88% effective at reducing the impact of head and brain injuries due to bicycle crashes.¹² Despite this, less than one quarter of bicyclists wear helmets.¹²
- In 2009, alcohol use was associated with nearly four in ten motor vehicle-related fatalities nationwide and in Vermont.¹³ Additionally, research examining drugs other than alcohol indicates that marijuana is the most prevalent drug detected in impaired drivers, fatally injured drivers, and motor vehicle crash victims nationwide.¹⁴

Alcohol, Tobacco, and Other Drugs

Early use of alcohol, tobacco, and other drugs is associated with myriad problems later in life. The high school and middle school surveys asked about tobacco, alcohol, marijuana, inhalant, and prescription drug use. The surveys also asked the age at which students first used alcohol, cigarettes, and marijuana. The high school survey asked additional questions on other drug use, such as heroin, methamphetamines, and hallucinogens. Both surveys asked about ease of availability of alcohol, cigarettes, and marijuana, peer disapproval of use of these substances, perceived parental disapproval of use, and the risk of harm associated with use of these substances.

- Alcohol use is a major contributing factor in about half of all homicides and sexual assaults,¹⁵ and about one-third of all motor vehicle crash fatalities.¹⁶ Heavy drinking among youth has been linked to violence, academic and job problems, suicidal behavior, trouble with law enforcement authorities, risky sexual behavior, and use of cigarettes, marijuana, cocaine, and other illegal drugs.^{17,18}
- Tobacco use is the single most preventable cause of death in the United States,¹⁹ contributing to more than one of every five deaths.²⁰ Cigarette smoking increases the risk of heart disease, chronic obstructive pulmonary disease, acute respiratory illness, stroke, and cancers of the lung, larynx, oral cavity, pharynx, pancreas, and cervix.¹⁹
- Marijuana use is associated with smoking-related respiratory damage, temporary short-term memory loss, decreased motivation, and psychological dependence.²¹ Other reactions include feelings of distrust, anxiety, or depression.²¹ In Vermont, more teens enter treatment with a primary diagnosis for marijuana dependence than all other illicit drugs combined.²²
- Other drug use is related to suicide, early unwanted pregnancy, school failure, delinquency, and transmission of sexually transmitted diseases (STDs), including HIV infection.²³
- Multi-year results from the Monitoring the Future survey indicate that the prevalence of marijuana use among youth declines as the percentage of youth expressing disapproval of marijuana increases.²⁴ As perception of harm of using alcohol and other drugs decreases, there is a tendency for use to increase.²⁴ Increased use is also associated with the perception that substances are readily available, regardless of whether the perception is accurate.²⁴

Sexual Behavior and Orientation

Experimentation with sexual behaviors and orientation often starts during adolescence. The high school survey asked about age of first sexual intercourse, frequency, sexual partners, sexual orientation, alcohol and drug use related to sexual intercourse, contraceptive use, and HIV testing. The middle school survey did not ask any questions about sex.

- Early sexual activity and having multiple sexual partners are associated with an increased risk of unwanted pregnancy, sexually transmitted diseases (STDs) including HIV infection,²⁵ and negative effects on social and psychological development.²⁶ Alcohol and drug use may serve as predisposing factors for initiation of sexual activity.²⁷
- Of the nearly 19 million new cases of STDs per year in the United States, almost half are among youth ages 15-24.²⁸ STDs may result in infertility and facilitation of HIV transmission and may have an adverse effect on pregnancy outcomes and maternal and child health.²⁶ Besides abstinence, condom use is currently the most effective means of preventing sexual transmission of HIV and other STDs.²⁹
- Although many lesbian, gay, bisexual, and transgender adolescents lead happy and healthy lives, others face tremendous challenges to growing up physically and mentally healthy. Compared to heterosexual youth, lesbian, gay, bisexual, and transgender youth are at higher risk for depression, tobacco, alcohol and other drug use, suicide, and unhealthy sexual behaviors.³⁰

Body Image

Negative feelings about weight and body image often develop in adolescence. The high school and middle school surveys included questions about weight perception and weight control. The high school survey asked for students' height and weight to calculate body mass index, and about use of artificial tanning devices.

- There are more than three times as many overweight children and adolescents in the U.S. than there were in 1980.³¹ Obesity in childhood and adolescence is associated with negative psychological and social consequences and adverse health outcomes, including type 2 diabetes, obstructive sleep apnea, hypertension, dyslipidemia, and metabolic syndrome.³² Overweight and obesity acquired during childhood or adolescence may persist into adulthood.³³ Approximately 400,000 deaths a year in the United States are currently associated with overweight and obesity.²⁰
- However, overemphasis on slenderness during adolescence may contribute to eating disorders such as anorexia nervosa and bulimia.^{34,35} Studies have shown high rates of body dissatisfaction and dieting among adolescent females, with many engaging in unhealthy weight control behaviors such as fasting and self-induced vomiting that can lead to abnormal physical and psychological development.^{36,37}
- Indoor tanning increases risk for melanoma and non-melanoma cancers. In the U.S., the incidence of melanoma is increasing more rapidly than that of any other cancer, particularly among girls and women between the ages of 15 and 39. Some evidence suggests that repeated UV irradiation, and the use of indoor tanning beds specifically, may have important behavioral consequences, including mood changes, pain, and physical dependency.³⁸

Nutrition and Physical Activity

Nutritious eating and physical activity are two cornerstones of healthy adolescent development. The high school and middle school surveys asked about breakfast consumption, physical activity, physical education classes, and television, computer, and video game use. The high school survey also asked about consumption of fruits, vegetables, soda, and sugar-sweetened beverages.

- Fruits and vegetables are good sources of complex carbohydrates, vitamins, minerals, and other substances that are important for good health. Dietary patterns with higher intakes of fruits and vegetables are associated with a variety of health benefits, including a decreased risk for some types of cancer.³⁹⁻⁴³
- In recent years, soft drink consumption has significantly increased among children and adolescents. Consumption of sugar-sweetened drinks, including soft drinks, appears to be associated with an increased risk for being overweight in children.⁴⁴⁻⁴⁵
- Regular physical activity builds and maintains healthy bones and muscles, controls weight, builds lean muscle, reduces fat, reduces feelings of depression and anxiety. It also decreases the risk of dying prematurely, dying of heart disease, and developing diabetes, colon cancer, and high blood pressure.⁴⁶ The U.S. Department of Health and Human Services recommends that young people ages 6–17 participate in at least 60 minutes of physical activity every day.⁴⁶
- By 12th grade, more than half of female students in the U.S. do not participate in vigorous physical activity regularly. School physical education classes can increase adolescent participation in physical activity and help adolescents develop the knowledge, attitudes, and skills they need to engage in lifelong physical activity.⁴⁷⁻⁵⁰
- Television viewing is the principal sedentary leisure time behavior in the U.S. Studies have shown that television viewing in young people is related to obesity⁵¹ and violent or aggressive behavior.⁵²⁻⁵⁴ Using the computer for fun and playing video games have become increasingly common sedentary leisure time activities among young people as well.

Measures of Youth Assets

Adolescent achievement requires sources of positive influence. The high school and middle school surveys asked about school and community connectivity as well as parental conversations about school. The high school survey asked additional questions about volunteerism and grades earned in school.

- Above-average school performance is one of many developmental assets, or factors promoting positive development. Studies have shown that students who get higher grades in school are less likely to use cigarettes, alcohol, or marijuana, and are more likely to postpone sexual intercourse.⁵⁵
- One of the strongest predictors of students' success in school is the extent to which their parents stay involved with their schoolwork.⁵⁶ A national study of adolescent health found that youth who reported a "connectedness" to their parents/family and school were the least likely to engage in risky behaviors.⁵⁷ Parental expectations regarding school achievement were also associated with lower levels of risk behaviors.⁵⁷
- Research shows that involvement in constructive, supervised extracurricular activities is associated with reduced likelihood of involvement in risky behaviors such as school failure, drug use, and delinquency.⁵⁸ In addition, evidence is emerging that students who participate in such activities are also more likely to engage in other "thriving" behaviors.⁵⁹
- Youth are not simply objects of adult efforts to modify their behaviors. Rather, if given the opportunities, they can make significant contributions to their families, schools, and communities. Adolescents, especially, need to exercise decision-making power in as many settings as is practical, so that they can develop into competent adults. Schools are a natural setting for youth to share in decisions that affect their lives.

References

1. Sosin, D.M., Koepsell, T.D., Rivara, F.P., Mercy, J.A. Fighting as a marker for multiple problem behaviors in adolescents. Journal of Adolescent Health 16(3):209-215, 1995.
2. Borowsky, I.W., Ireland, M. Predictors of future fight-related injury among adolescents. Pediatrics 113(3 pt 1):530-536, 2005.
3. Pickett, W., Craig, W., Harel, Y., et al. Cross-national study of fighting and weapon carrying as determinants of adolescent injury. Pediatrics 116(6):e855-863, 2005.
4. Roberts, T.A., Klein, J.D., Fisher, S. Longitudinal effect of intimate partner abuse and high-risk behavior among adolescents. Archives of Pediatrics and Adolescent Medicine 157(9):875-881, 2003.
5. Ackard, D.M., Neumark-Sztainer, D. Date violence and date rape among adolescents: association with disordered eating behaviors and psychological health. Child Abuse and Neglect 26(5):455-473, 2002.
6. Howard, D.E., Wang, M.Q. Psychosocial correlates of U.S. adolescents who report a history of forced sexual intercourse. Journal of Adolescent Health 36(5):372-379, 2005.
7. Juvonen, J., Graham, S., Schuster, M.A. Bullying among young adolescents: the strong, the weak, and the troubled. Pediatrics 112(6 pt 1): 1231-1237, 2003.
8. Spivak, H., Prothrow-Stith, D. The need to address bullying-an important component of violence prevention. JAMA 285(16):2131-2132, 2001.
9. Nansel, T.R., Overpeck, M., Pilla, R.S., et al. Bullying behaviors among U.S. youth: prevalence and association with psychological adjustment. JAMA 285(16):2094-2100, 2001.
10. Centers for Disease Control and Prevention. Web-based Injury Statistics Query and Reporting System (WISQARS). National Center for Injury Prevention and Control, Centers for Disease Control and Prevention. Online: www.cdc.gov/injury/wisqars/index.html
11. National Highway Traffic Safety Administration. Traffic safety facts: occupant protection, 2007. Online: www-nrd.nhtsa.dot.gov/Pubs/810991.PDF
12. National Highway Traffic Safety Administration. Traffic safety facts: bicycle helmet use laws, 2008. Online: www.nhtsa.dot.gov/people/injury/TSFLaws/PDFs/810886.pdf
13. National Highway Traffic Safety Administration. Traffic safety facts: state alcohol-impaired driving estimates, 2009. Online: www-nrd.nhtsa.dot.gov/Pubs/811398.pdf
14. Jones, R.K., Shinar, D., Walsh, J.M. State of Knowledge of Drug-Impaired Driving. National Highway Traffic Safety Administration Technical Report DOT HS 809 642. Washington, DC: U.S. Department of Transportation, 2003.
15. Abbey, A., Zawacki, T., Buck, P.O., et al. Alcohol and sexual assault. Alcohol Research and Health 25(1):43-51, 2001.
16. Miller, J.W., Naimi, T.S., Brewer, R.D., Jones, S.E. Binge drinking and associated health risk behaviors among high school students. Pediatrics 119(1):76-85, 2007.

References

17. National Research Council and Institute of Medicine (2004). Reducing Underage Drinking: A Collective Responsibility. Committee on Developing a Strategy to Reduce and Prevent Underage Drinking, Richard J. Bonnie and Mary Ellen O'Connell, Editors. Board on Children, Youth, and Families, Division of Behavioral and Social Sciences and Education. Washington, DC: The National Academies Press.
18. U.S. Department of Health and Human Services. The Surgeon General's Call to Action to Prevent and Reduce Underage Drinking. U.S. Department of Health and Human Services, Office of the Surgeon General, 2007.
19. U.S. Department of Health and Human Services. The Health Consequences of Smoking: A Report of the Surgeon General. U.S. Department of Health and Human Services; Centers for Disease Control and Prevention; National Center for Chronic Disease Prevention and Health Promotion: Office on Smoking and Health, 2004.
20. Mokdad, A.H., Marks, J.S., Stroup, D.F., Gerberding, J.L. Actual causes of death in the United States, 2000. JAMA 291(10):1238-1245, 2004.
21. National Institute on Drug Abuse. Research Report Series: Marijuana Abuse (NIH Publication 05-3859). Bethesda, MD: National Institute on Drug Abuse, 2002.
22. Vermont Substance Abuse Treatment Information System. Data online at: www.healthvermont.gov/adap/clearinghouse/documents/AdolescentsbySAandFY.pdf
23. Newcomb, M.D., Locke T. Health, social, and psychological consequences of drug use and abuse. In: Epidemiology of Drug Abuse (Z. Sloboda, ed.). Springer U.S., 2006.
24. Johnston, L., O'Malley, P., Bachman, J. G., Shulenberg, J.E. National Survey Results on Drug Use From the Monitoring the Future Study, 1975-2007, Volume I: Secondary School Students (NIH Publication No. 08-6418A). Bethesda, MD: National Institute of Drug Abuse, 2008.
25. Abma JC, Martinez GM, Copen CE. Teenagers in the United States: Sexual activity, contraceptive use, and childbearing, National Survey of Family Growth 2006-2008. National Center for Health Statistics. Vital Health Stat 23 (30). 2010.
26. Centers for Disease Control and Prevention. Fact sheet on STDs and pregnancy. Online: www.cdc.gov/std/pregnancy/STDs-and-pregnancy-fact-sheet.pdf
27. Cavazos-Rehg, P.A., Krauss, M.J., Spitznagel, E.L., et al. Substance use and the risk for sexual intercourse with and without a history of teenage pregnancy among adolescent females. Journal of Studies on Alcohol and Drugs 72(2): 194-198, 2011.
28. Gavin, L., MacKay, A.P., Brown, K., et al. Sexual and reproductive health of persons aged 10-24 years – United States, 2002-2007. MMWR Surveillance Summaries 58(6): 1-58, 2009.
29. Joint United Nations Programme on HIV/AIDS (UNAIDS). Fast Facts about HIV Prevention. Online at: www.unaids.org/en/media/unaids/contentassets/dataimport/pub/basedocument/2008/20080501_fastfacts_prevention_en.pdf
30. Kann, L., Olsen, E.O., McManus, T., et al. Sexual Identity, Sex of Sexual Contacts, and Health-Risk Behaviors Among Students in Grades 9-12 — Youth Risk Behavior Surveillance, Selected Sites, United States, 2001-2009. MMWR Early Release 60(7): 1-133, 2011.
31. Ogden, C.L., Carroll, M.D. Prevalence of obesity among children and adolescents: United States, trends 1963-1965 through 2007-2008. National Center for Health Statistics Health E-Stats, June 2010.

References

32. Daniels, S.R., Arnett, D.K., Eckel, R.H., et al. Overweight in children and adolescents: pathophysiology, consequences, prevention, and treatment. Circulation 111(15):1999-2012, 2005.
33. Wright, C.M., Parker, L., Lamont, D., Craft, A.W. Implications of childhood obesity for adult health: findings from thousand families cohort study. British Medical Journal 323(7324):1280-1284, 2001.
34. Tremblay, L., Lariviere, M. The influence of puberty onset, body mass index, and pressure to be thin on disordered eating behaviors in children and adolescents. Eating Behaviors 10(2):75-83, 2009.
35. Mitchell, J.E., Eckert, E.D. Scope and significance of eating disorders. Journal of Consulting Clinical Psychology 55:628-634, 1987.
36. Neumark-Sztainer, D., Hannan, P.J. Weight-related behaviors among adolescent girls and boys: results from a national survey. Archives of Pediatric and Adolescent Medicine 154(6):569-577, 2000.
37. Neumark-Sztainer, D., Story, M., Hannan, P.J., et al. Weight-related concerns and behaviors among overweight and nonoverweight adolescents: implications for preventing weight-related disorders. Archives of Pediatric and Adolescent Medicine 156(2):171-178, 2002.
38. Fisher, D.E. and James, W.D. Indoor tanning — science, behavior, and policy. New England Journal of Medicine 363:901-903, 2010.
39. Key, T.J., Schatzkin, A., Willet, W.C., et al. Diet, nutrition, and the prevention of cancer. Public Health Nutrition 7(1A):187-200, 2004.
40. National Cancer Institute. 5 A Day for Better Health Program (NIH Publication 01-5019). Bethesda, MD, 2001.
41. Kavey, R.E., Daniels, S.R., Lauer, R.M., et al. American Heart Association guidelines for primary prevention of atherosclerotic cardiovascular disease beginning in childhood. Journal of Pediatrics 142(4):368-372, 2003.
42. Terry, P., Terry, J.B., Wolk, A. Fruit and vegetable consumption in the prevention of cancer: an update. Journal of Internal Medicine 250(4):280-290, 2001.
43. Van Duyn, M.A., Pivonka, E. Overview of the health benefits of fruit and vegetable consumption for the dietetics professional: selected literature. Journal of the American Dieticians Association 100(12):1511-1521, 2000.
44. Malik, V.S., Schulze, M.B., Hu, F.B. Intake of sugar-sweetened beverages and weight gain: a systematic review. American Journal of Clinical Nutrition 84(2):274-288, 2006.
45. Ludwig, D.S., Peterson, K.E., Gortmaker, S.L. Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. Lancet 357(9255):505-508, 2001.
46. Physical Activity Guidelines Advisory Committee. Physical Activity Guidelines Advisory Committee Report, 2008. Washington, D.C.: U.S. Department of Health and Human Services, 2008.
47. Durant, N., Harris, S.K., Doyle, S., et al. Relation of school environment and policy to adolescent physical activity. Journal of School Health 79(4):153-159, 2009.
48. McKenzie, K.L., Li, D., Derby, C.A., et al. Maintenance of effects of the CATCH Physical Education Program: results from the CATCH-ON Study. Health Education & Behavior 30(4):447-462, 2003.

References

49. U.S. Department of Health and Human Services and U.S. Department of Education. Promoting better health for young people through physical activity and sports. 2000. Online: www.cdc.gov/HealthyYouth/physicalactivity/promoting_health/pdfs/ppar.pdf
50. Center for Disease Control and Prevention. Guidelines for school and community programs to promote lifelong physical activity among young people. Morbidity and Mortality Weekly Report 46 (No. RR-6):1-36, 1997.
51. Zabinski, M.F., Norman, G.J., Sallis, J.F., et al. Patterns of sedentary behavior among adolescents. Health Psychology 26(1):113-120, 2007.
52. Crespo, C.J., Smit, E., Troiano, R.P., et al. Television watching, energy intake, and obesity in U.S. children: results from the third National Health and Nutrition Examination Survey, 1988-1994. Archives of Pediatric and Adolescent Medicine 155(3):360-365, 2001.
53. Kaur, H., Choi, W.S., Mayo, M.S., Harris, K.J. Duration of television watching is associated with increased body mass index. Journal of Pediatrics 143(4):506-511, 2003.
54. Kuntsche, E., Pickett, W., Overpeck, M., et al. Television viewing and forms of bullying among adolescents from eight countries. Journal of Adolescent Health 39(6):908-915, 2006.
55. Resnick, M.D., Bearman, P.S., Blum, R.W., et al. Protecting adolescents from harm. Findings from the National Longitudinal Study on Adolescent Health. JAMA 278(10):823-832, 1997.
56. Fan, X., Chen, M. Parental involvement and students' academic achievement: a meta-analysis. Educational Psychology Review 13(1):1-22, 2001.
57. U.S. Council of Economic Advisors. Teens and their parents in the 21st century: An examination of trends in teen behavior and the role of parental involvement. 2000. Online: http://clinton3.nara.gov/WH/EOP/CEA/html/Teens_Paper_Final.pdf
58. Fredricks, J.A., Eccles, J.S. Is extracurricular participation associated with beneficial outcomes? Developmental Psychology 42(4):698-713, 2006.
59. Scales, P.C., Benson, P.L., Leffert, N., Blyth, D.A. Contribution of developmental assets to prediction of thriving among adolescents. Applied Developmental Science 4(1):27-46, 2000

2011 Vermont High School Youth Risk Behavior Survey

Results for Washington South SU

Participation Rates

	Grade				Gender		All
	9	10	11	12	F	M	
Participated	46	44	40	32	89	73	165
Enrolled	54	50	55	40	98	101	199
Participation Rate	85%	88%	71%	80%	90%	72%	82%

Based on October 1, 2010 enrollment. Some students did not indicate grade or gender.

Demographics

Age

	2011
14 or younger	18%
15	21%
16	26%
17	23%
18 or older	12%

Race and Ethnicity

	2011
White non-Hispanic	87%
Racial or ethnic minority	13%

Mother's Education

	2011
High school or less	39%
Some college	13%
College graduate	30%
Graduate or professional degree after college	11%
Not sure	7%

Personal Safety

Percent of students who:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Ride bicycles	69%	64%	84%	63%	63%	61%	79%

Of students who ride bicycles, percent who wear helmets:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Always	12%	*	22%	*	*	*	18%
Most of the time	8%	*	*	*	*	11%	*
Sometimes	10%	18%	*	*	*	17%	*
Rarely or never	70%	68%	62%	75%	80%	67%	73%

* Suppressed due to small cell sizes (≤ 5 students)

Personal Safety

Percent of students who wear seatbelts when riding in a car:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Always	53%	45%	61%	50%	56%	50%	57%
Most of the time	24%	30%	20%	24%	22%	30%	17%
Sometimes	11%	16%	*	*	*	11%	11%
Rarely or never	11%	*	*	16%	*	9%	14%

Percent of students who, in the past 30 days:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Rode in a car or other vehicle driven by someone who had been drinking alcohol	28%	20%	23%	34%	38%	30%	25%
Drove a car or other vehicle after drinking	9%	*	*	*	28%	8%	11%
Rode in a car or other vehicle driven by someone who had been smoking marijuana	30%	13%	14%	39%	65%	30%	30%
Drove a car or other vehicle after smoking marijuana	15%	*	*	22%	34%	13%	17%

* Suppressed due to small cell sizes (≤ 5 students)

Personal Safety

Percent of students who:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Carried a weapon such as a gun, knife or club on school property, past 30 days	10%	*	*	*	22%	7%	14%
Did not go to school because they felt unsafe, past 30 days	4%	*	*	*	*	7%	*
Were threatened or injured with a weapon at school, past 12 months	5%	*	*	*	*	*	*
Were in a physical fight, past 12 months	33%	36%	27%	39%	28%	20%	48%
Were in a physical fight on school property, past 12 months	16%	18%	18%	18%	*	7%	27%
Were hit, slapped, or physically hurt on purpose by a boyfriend or girlfriend, past 12 months	9%	*	*	*	*	11%	*
Were physically forced to have sexual intercourse, ever	6%	*	*	*	*	8%	*

* Suppressed due to small cell sizes (≤ 5 students)

Personal Safety

Percent of students who:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Were bullied, past 30 days	22%	24%	23%	24%	*	25%	19%
Bullied someone else, past 30 days	22%	27%	19%	16%	28%	19%	26%
Were electronically bullied, past 12 months	21%	16%	23%	18%	28%	28%	11%
Caused purposeful harm to themselves such as by cutting or burning, past 12 months	16%	16%	14%	24%	*	21%	10%
Felt sad or hopeless almost every day for two weeks or more in a row, past 12 months	30%	33%	30%	29%	25%	39%	19%
Made a suicide plan, past 12 months	15%	22%	20%	*	*	17%	13%
Made a suicide attempt, past 12 months	4%	*	*	*	*	*	*

For the purposes of the Vermont YRBS, bullying was described as occurring when, on many occasions, a student or group of students say or do unpleasant things to another student to make fun of, tease, embarrass, or scare him/her, or purposefully exclude him/her. Bullying can occur before, during, or after the school day; on school property; on a school bus; or at a school-sponsored activity. It is not bullying when two students of about the same strength and power argue or fight, or when teasing is done in a friendly way. Electronic bullying includes through e-mail, chat rooms, instant messaging, Web sites, or texting.

* Suppressed due to small cell sizes (≤ 5 students)

Alcohol, Tobacco, and Other Drug Use

Percent of students who:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Ever drank alcohol other than a few sips	69%	56%	67%	69%	91%	70%	69%
Had first drink other than a few sips before age 13	21%	22%	23%	17%	19%	22%	19%
Drank alcohol, past 30 days	34%	22%	19%	36%	69%	30%	40%
Had five or more drinks of alcohol in a row, past 30 days	23%	*	14%	31%	53%	20%	28%
Drank alcohol on 3 to 9 of the past 30 days	13%	*	*	19%	31%	10%	16%
Drank alcohol on 10 or more of the past 30 days	5%	*	*	*	*	6%	*
Binged on 3 or more of the past 30 days	10%	*	*	*	25%	9%	10%
Drank alcohol on school property, past 30 days	5%	*	*	*	*	*	*

* Suppressed due to small cell sizes (≤ 5 students)

Alcohol, Tobacco, and Other Drug Use

Among students who drank in the past 30 days:

Usual type of beverage consumed, past 30 days

	All	Sex	
	2011	Female	Male
Liquor, such as rum, gin, vodka, or whiskey	43%	54%	33%
Beer	29%	*	48%
Other alcoholic beverages, such as Smirnoff Ice, Bacardi Silver, Mike's Hard Lemonade, or Twisted Tea	22%	33%	*
Wine	*	*	*

Usual source of alcohol, past 30 days

	All	Sex	
	2011	Female	Male
Someone gave it to me	35%	35%	35%
Gave someone money to buy it for me	46%	46%	46%
Got it or stole it from home	13%	19%	*
Purchased from a store, liquor store; or restaurant	*	*	*
Stole it from a store or restaurant	*	*	*

* Suppressed due to small cell sizes (≤ 5 students)

There were not enough current drinkers to report usual source or type of alcohol by grade.

Alcohol, Tobacco, and Other Drug Use

Percent of students who:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Ever smoked a whole cigarette	37%	20%	23%	46%	69%	37%	37%
Smoked before age 13	13%	*	*	24%	*	8%	20%
Smoked cigarettes, past 30 days	26%	18%	*	37%	50%	25%	28%
Smoked on 20 or more of the past 30 days	12%	*	*	*	28%	8%	17%
Smoked every day, past 30 days	9%	*	*	*	19%	6%	13%
Used chewing tobacco, past 30 days	8%	*	*	*	*	*	13%
Smoked cigars, past 30 days	13%	*	*	24%	22%	10%	15%
Used snus, ever	11%	*	*	*	25%	11%	11%

* Suppressed due to small cell sizes (≤ 5 students)

Alcohol, Tobacco, and Other Drug Use

Among students who smoked in the past 30 days:

Usual source of cigarettes, past 30 days

	All	Sex	
	2011	Female	Male
Bought them in a store or from a vending machine	24%	*	35%
Gave someone else money to buy them for me	22%	*	30%
Borrowed (or bummed) them from someone else; Person 18 years or older gave them to me; took them from a store or family member	41%	67%	*
Some other way	*	*	*

Percent who tried to quit, past 12 months

	All	Sex	
	2011	Female	Male
Tried to quit	44%	38%	50%

* Suppressed due to small cell sizes (≤ 5 students)

There were not enough current smokers to report usual source of cigarettes or quit attempts by grade.

Alcohol, Tobacco, and Other Drug Use

Percent of students who:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Were in the same room with someone who was smoking, past 7 days	55%	44%	41%	61%	84%	56%	55%
Were in the same car with someone who was smoking, past 7 days	50%	44%	34%	55%	72%	50%	49%
Were asked about smoking by a health professional, past 12 months	35%	27%	36%	29%	53%	42%	27%
Discussed dangers of smoking with parents or guardian, past 12 months	35%	29%	34%	49%	31%	32%	39%
Think that out of 100 Vermont high school students, 56 or more smoke cigarettes	21%	16%	26%	29%	13%	23%	19%

Alcohol, Tobacco, and Other Drug Use

Percent of students who:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Used marijuana, ever	44%	27%	34%	53%	74%	41%	48%
Used marijuana before age 13	9%	*	*	*	*	7%	11%
Used marijuana, past 30 days	29%	16%	18%	33%	56%	27%	30%
Used marijuana on 3 to 9 of the past 30 days	*	*	*	*	*	*	*
Used marijuana on 10 or more of the past 30 days	18%	*	*	28%	38%	18%	19%
Used marijuana on school property, past 30 days	8%	*	*	*	*	9%	*

* Suppressed due to small cell sizes (≤ 5 students)

Alcohol, Tobacco, and Other Drug Use

Percent of students who:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Used cocaine, past 30 days	9%	*	*	*	*	11%	*
Used inhalants, ever	13%	*	*	21%	*	14%	11%
Used heroin, ever	7%	*	*	*	*	7%	*
Used methamphetamines, ever	8%	*	*	*	*	8%	*
Used hallucinogens, ever	18%	*	*	29%	34%	15%	21%
Used the prescription stimulant Ritalin or Adderall not prescribed to them, ever	12%	*	*	22%	*	14%	10%
Used a prescription pain reliever such as OxyContin, Vicodin, or other prescription pain reliever not prescribed to them, ever	20%	*	14%	34%	31%	19%	21%
Used a prescription pain reliever or a prescription stimulant not prescribed to them, ever	22%	*	16%	39%	31%	22%	23%
Were offered, sold, or given an illegal drug on school property, past 12 months	21%	*	*	37%	28%	19%	23%

* Suppressed due to small cell sizes (≤ 5 students)

Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use

Percent of students who think their parents would think it is wrong or very wrong for them to:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Smoke cigarettes	84%	93%	89%	84%	63%	85%	82%
Drink alcohol	76%	89%	80%	74%	56%	81%	70%
Smoke marijuana	84%	91%	93%	74%	75%	83%	86%

Percent of students who think it would be wrong or very wrong for someone their age to:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Smoke cigarettes	72%	89%	84%	66%	41%	70%	75%
Drink alcohol	54%	76%	58%	45%	31%	56%	52%
Smoke marijuana	63%	80%	73%	47%	44%	63%	63%

Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use

Percent of students who think people their age greatly risk harming themselves (physically or in other ways) if they:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Smoke one or more packs of cigarettes per day	61%	73%	58%	59%	47%	60%	61%
Have five or more drinks of alcohol once or twice each weekend	35%	44%	40%	32%	19%	36%	34%
Smoke marijuana regularly	39%	66%	40%	28%	13%	45%	32%

Percent of students who report that it would be easy or very easy to get:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Cigarettes	69%	56%	60%	76%	91%	71%	66%
Alcohol	69%	58%	69%	82%	71%	70%	69%
Marijuana	62%	47%	45%	79%	88%	66%	59%

Sexual Behavior and Orientation

Percent of students who:

	All	Grade		Sex	
	2011	9 and 10	11 and 12	Female	Male
Had sexual intercourse, ever	49%	29%	76%	50%	48%
First had sexual intercourse before age 13	6%	7%	*	*	*
Had sexual intercourse with four or more people during their life	21%	11%	33%	22%	18%
Had sexual intercourse at least once in the past three months	40%	25%	60%	41%	39%
Had oral sex, ever	52%	33%	76%	57%	45%
Had a HIV test, ever	12%	8%	18%	20%	*

* Suppressed due to small cell sizes (≤ 5 students)

Sexual Behavior and Orientation

Among students who had sex in the past 3 months:

Percent of students who:

	All	Grade		Sex	
	2011	9 and 10	11 and 12	Female	Male
Used drugs or alcohol before last sexual intercourse	33%	27%	37%	34%	32%
Used a condom for last sexual intercourse	60%	59%	61%	49%	75%

Method used to prevent pregnancy during last sexual intercourse

	All	Grade		Sex	
	2011	9 and 10	11 and 12	Female	Male
Condoms	41%	60%	32%	32%	52%
Prescription birth control including pills, rings, implants, injections, IUDs	41%	*	51%	47%	33%
No method	*	*	*	14%	*
Withdrawal	8%	*	*	*	*
Some other method/not sure	*	*	*	*	*

* Suppressed due to small cell sizes (≤ 5 students)

Sexual Behavior and Orientation

Sexual orientation

	Sex	
	Female	Male
Heterosexual	80%	96%
Gay or lesbian	*	*
Bisexual	13%	*
Not sure	*	*

With whom have you had sexual contact?

	Sex	
	Female	Male
Females	*	61%
Males	44%	*
Both	15%	*
Never had sexual contact	39%	39%

* Suppressed due to small cell sizes (≤ 5 students)

Body Image

Percent of students who are:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Overweight (85th BMI Percentile)	21%	26%	23%	23%	*	22%	20%
Obese (95th BMI Percentile)	10%	*	*	*	*	8%	12%

BMI=body mass index; weight in kilograms divided by height in meters, squared

Percent of students who think that they are:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Underweight	12%	13%	*	*	19%	7%	18%
About the right weight	53%	51%	59%	53%	50%	51%	56%
Overweight	35%	36%	34%	37%	31%	42%	25%

* Suppressed due to small cell sizes (≤ 5 students)

Body Image

Percent of students who are trying to:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Lose weight	47%	52%	49%	43%	40%	61%	29%
Gain weight	12%	14%	19%	*	*	*	25%
Stay the same weight	25%	20%	23%	32%	27%	24%	28%
Not trying to do anything about weight	16%	14%	*	19%	23%	13%	19%

Percent of students who:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Skipped meals to lose weight or keep from gaining weight, past 30 days	19%	23%	18%	16%	19%	30%	*
Used an indoor tanning device such as a sunlamp, sunbed, or tanning booth, past 30 days	9%	*	*	16%	*	14%	*

* Suppressed due to small cell sizes (≤ 5 students)

Nutrition

Percent of students who, during the past seven days:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Ate fruits and vegetables five or more times per day	23%	22%	19%	32%	22%	23%	24%
Ate two or more servings of fruit or fruit juice per day	32%	31%	34%	32%	31%	34%	30%
Ate three or more servings of vegetables per day	19%	18%	16%	26%	16%	19%	19%
Ate breakfast seven of the past seven days	42%	38%	50%	42%	38%	42%	42%
Ate breakfast at least three of the past seven days	75%	80%	73%	68%	81%	72%	80%
Ate breakfast zero of the past seven days	11%	*	*	*	*	10%	13%
Drank a can, bottle, or glass of soda or pop at least once per day	20%	18%	20%	18%	25%	10%	32%
Drank a can, bottle, or glass of a sugar-sweetened beverage such as lemonade, sweetened tea or coffee drinks, sports drinks, energy drinks, Snapple, or Sunny Delight at least once per day	22%	22%	18%	18%	31%	18%	27%

* Suppressed due to small cell sizes (≤ 5 students)

Physical Activity

Percent of students who were physically active on:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Seven of the past seven days	31%	27%	41%	26%	28%	24%	39%
Zero of the past seven days	16%	20%	*	26%	*	18%	14%

Percent of students who attend physical education classes:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Every day during an average school week	48%	69%	50%	42%	23%	55%	40%
At least one day during an average school week	50%	71%	51%	46%	25%	56%	43%

Percent of students who, on an average school day, watch TV, play video or computer games, or use a computer for something that is not school work:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Three or more hours	42%	47%	32%	46%	44%	41%	42%
Five or more hours	18%	24%	16%	16%	*	17%	18%

* Suppressed due to small cell sizes (≤ 5 students)

Youth Assets

Grades earned in school

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Mostly As	26%	41%	19%	26%	*	37%	12%
Mostly Bs	45%	25%	53%	47%	60%	43%	47%
Mostly Cs	20%	18%	14%	26%	23%	12%	29%
Mostly Ds or Fs	8%	16%	*	*	*	*	12%
None/not sure	*	*	*	*	*	*	*

Percent of students who talk with their parents about school:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
About every day	45%	40%	48%	49%	44%	45%	44%
About once or twice a week	28%	36%	23%	20%	34%	26%	31%
About once or twice a month	8%	*	*	*	*	10%	*
Less than once a month	11%	*	*	*	*	11%	11%
Never	8%	*	*	*	*	9%	*

* Suppressed due to small cell sizes (≤ 5 students)

Youth Assets

Percent of students who:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Volunteer one or more hours in an average week	55%	51%	44%	62%	66%	50%	61%
Volunteer three or more hours in an average week	15%	13%	14%	16%	16%	13%	17%

Percent of students who agree or strongly agree:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Teachers really care about you and give you a lot of encouragement	59%	49%	63%	59%	66%	57%	61%
Students help decide what goes on in your school	40%	42%	52%	27%	34%	45%	33%
In your community, you feel like you matter to people	45%	42%	53%	43%	41%	45%	45%

2011 Vermont Middle School Youth Risk Behavior Survey

Results for Washington South SU

Participation Rates

	Grade			Gender		All
	6	7	8	F	M	
Participated	34	45	43	59	63	122
Enrolled	39	49	48	63	73	136
Participation Rate	87%	92%	90%	94%	86%	90%

Based on October 1, 2010 enrollment.

Demographics

Age

	All
	2011
11 or younger	16%
12	30%
13	34%
14 or older	20%

Race and Ethnicity

	All
	2011
White non-Hispanic	84%
Racial or ethnic minority	16%

Personal Safety

Percent of students who:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Ride bicycles	95%	100%	98%	87%	94%	95%

Of students who ride bicycles, percent who wear helmets:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Always	22%	30%	21%	*	29%	16%
Most of the time	17%	*	24%	*	12%	21%
Sometimes	17%	21%	14%	*	18%	16%
Rarely or never	44%	33%	40%	61%	41%	47%

* Suppressed due to small cell sizes (≤ 5 students)

Personal Safety

Percent of students who:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Rollerblade or skateboard	58%	65%	58%	53%	56%	60%

Of students who rollerblade or skateboard, percent who wear helmets:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Always	27%	36%	31%	*	39%	16%
Most of the time	*	*	*	*	*	*
Sometimes	13%	*	*	*	*	16%
Rarely or never	54%	41%	58%	61%	48%	58%

* Suppressed due to small cell sizes (≤ 5 students)

Personal Safety

Percent of students who wear seatbelts when riding in a car:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Always	66%	74%	64%	63%	68%	65%
Most of the time	28%	26%	27%	30%	25%	30%
Sometimes	*	*	*	*	*	*
Rarely or never	*	*	*	*	*	*

Percent of students who ever rode in a car driven by someone who had been drinking alcohol:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Yes	27%	21%	31%	28%	24%	30%
No	53%	62%	58%	42%	53%	54%
Not sure	27%	22%	*	42%	31%	23%

* Suppressed due to small cell sizes (≤ 5 students)

Personal Safety

Percent of students who:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Were ever in a physical fight	56%	41%	67%	56%	41%	70%
Did not go to school because felt unsafe, past 30 days	7%	*	*	*	10%	3%
Were ever bullied on school property	71%	59%	76%	77%	63%	79%
Were ever electronically bullied	35%	32%	25%	47%	47%	23%
Were bullied, past 30 days	42%	44%	29%	53%	53%	32%
Bullied someone else, past 30 days	31%	24%	27%	40%	29%	32%
Ever seriously thought about killing themselves	23%	*	24%	29%	32%	15%
Ever made a suicide plan	18%	18%	17%	19%	25%	12%
Ever attempted suicide	6%	*	*	*	*	*

* Suppressed due to small cell sizes (≤ 5 students)

Alcohol, Tobacco, and Other Drug Use

Percent of students who:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Ever had a drink of alcohol other than a few sips	36%	29%	45%	32%	40%	32%
Had first drink other than a few sips before age 11	17%	25%	14%	*	18%	15%
Drank alcohol, past 30 days	12%	*	20%	*	14%	10%
Had five or more drinks of alcohol in a row, past 30 days	*	*	*	*	*	*

* Suppressed due to small cell sizes (≤ 5 students)

Alcohol, Tobacco, and Other Drug Use

Percent of students who:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Tried cigarette smoking, even one or two puffs, ever	16%	*	18%	19%	19%	14%
Smoked before age 11	*	*	*	*	*	*
Smoked in past 30 days	*	*	*	*	*	*
Used chewing tobacco, past 30 days	*	*	*	*	*	*
Smoked cigars, past 30 days	*	*	*	*	*	*
Used snus, ever	6%	*	*	*	*	*

* Suppressed due to small cell sizes (≤ 5 students)

Alcohol, Tobacco, and Other Drug Use

Percent of students who:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Were in the same room with someone who was smoking, past 7 days	41%	39%	45%	37%	39%	43%
Were in the same car with someone who was smoking, past 7 days	25%	18%	31%	23%	24%	25%
Were asked about smoking by a health professional, past 12 months	17%	*	18%	21%	12%	22%
Discussed dangers of smoking with parents or guardian, past 12 months	51%	62%	51%	42%	51%	51%
Think that out of 100 Vermont high school students, 56 or more smoke cigarettes	18%	12%	20%	21%	20%	16%

* Suppressed due to small cell sizes (≤ 5 students)

Alcohol, Tobacco, and Other Drug Use

Percent of students who:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Used marijuana, ever	8%	*	*	*	*	*
Used marijuana before age 11	*	*	*	*	*	*
Used marijuana, past 30 days	*	*	*	*	*	*
Used inhalants, ever	11%	*	*	*	*	*
Misused prescription drugs, ever	*	*	*	*	*	*

* Suppressed due to small cell sizes (≤ 5 students)

Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use

Percent of students who think their parents would think it is wrong or very wrong for them to:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Smoke cigarettes	95%	97%	93%	95%	92%	98%
Drink alcohol	85%	88%	84%	84%	81%	89%
Smoke marijuana	95%	100%	91%	95%	95%	95%

Percent of students who think it would be wrong or very wrong for someone their age to:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Smoke cigarettes	89%	100%	84%	84%	86%	90%
Drink alcohol	76%	91%	73%	67%	75%	77%
Smoke marijuana	92%	100%	89%	88%	90%	94%

Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use

Percent of students who think people their age greatly risk harming themselves (physically or in other ways) if they:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Smoke one or more packs of cigarettes per day	65%	65%	67%	63%	66%	63%
Have five or more drinks of alcohol once or twice each weekend	34%	41%	27%	35%	29%	38%
Smoke marijuana regularly	62%	75%	53%	60%	62%	61%

Percent of students who report that it would be easy or very easy to get:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Cigarettes	38%	22%	32%	57%	37%	39%
Alcohol	44%	22%	50%	55%	44%	44%
Marijuana	10%	*	16%	10%	8%	12%

* Suppressed due to small cell sizes (≤ 5 students)

Body Image

Percent of students who think that they are:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Underweight	19%	21%	16%	21%	17%	21%
About the right weight	58%	62%	52%	60%	63%	53%
Overweight	23%	18%	32%	19%	20%	26%

Percent of students who are trying to:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Lose weight	48%	44%	47%	51%	54%	41%
Gain weight	11%	*	*	*	10%	13%
Stay the same weight	20%	24%	18%	21%	14%	27%
Not trying to do anything about weight	20%	21%	24%	16%	22%	19%

Percent of students who:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Skipped meals to lose weight or keep from gaining weight, past 30 days	18%	*	22%	23%	20%	16%

* Suppressed due to small cell sizes (≤ 5 students)

Nutrition

Percent of students who ate breakfast on:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Seven of the past seven days	47%	62%	44%	37%	42%	51%
At least three of the past seven days	80%	91%	78%	72%	71%	87%
Zero of the past seven days	7%	*	*	*	*	*

* Suppressed due to small cell sizes (≤ 5 students)

Physical Activity

Percent of students who were physically active on:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Seven of the past seven days	36%	38%	30%	40%	32%	39%
Zero of the past seven days	7%	*	*	*	*	*

Percent of students who attend physical education classes:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Every day during an average school week	*	*	*	*	*	*
At least one day during an average school week	98%	97%	100%	98%	98%	98%

* Suppressed due to small cell sizes (≤ 5 students)

Physical Activity

Percent of students who, on an average school day, watch TV:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Three or more hours	25%	29%	24%	21%	29%	21%
Five or more hours	6%	*	*	*	*	*

Percent of students who, on an average school day, play video or computer games or use a computer for something that is not school work:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Three or more hours	25%	*	24%	37%	24%	27%
Five or more hours	7%	*	*	*	*	10%

* Suppressed due to small cell sizes (≤ 5 students)

Youth Assets

Percent of students who, in the past 7 days, went online for something that was not for school:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Zero or one day	24%	38%	22%	14%	20%	27%
Two or three days	23%	26%	20%	23%	25%	21%
Four or five days	23%	29%	22%	19%	25%	21%
Six or seven days	30%	*	36%	44%	29%	32%

Percent of students who talk with their parents about school:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
About every day	45%	47%	53%	35%	47%	43%
About once or twice a week	22%	24%	18%	26%	15%	29%
About once or twice a month	15%	*	*	19%	17%	13%
Less than once a month	10%	*	*	16%	14%	*
Never	8%	*	13%	*	*	10%

* Suppressed due to small cell sizes (≤ 5 students)

Youth Assets

Percent of students who agree or strongly agree:

	All	Grade			Sex	
	2011	6	7	8	Female	Male
Teachers really care about you and give you a lot of encouragement	67%	76%	76%	51%	68%	67%
Students help decide what goes on in your school	47%	44%	51%	44%	46%	48%
In your community, you feel like you matter to people	52%	55%	49%	53%	48%	56%

Vermont High School Youth Risk Behavior Survey

Recalculated Results for Washington South SU

Before 2011, students in grades 8-12 took the Vermont Youth Risk Behavior Survey. In 2011, we conducted a high school survey of students in grades 9-12, and a middle school survey of students in grades 6-8.

All results in the 2011 high school survey report are for grades 9-12 only. The following pages contain rates from previous surveys recalculated for grades 9-12, overall and by gender, for comparison to 2011. Rates by grade did not change.

This section of the report includes only the variables on the 2011 YRBS. # = question was not asked that year.

Personal Safety - All

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Always/almost always wear seat belt when riding in car	69%	71%	77%	80%	71%	76%
Rode with drinking driver, past 30 days	32%	20%	17%	24%	22%	22%
Drove after drinking, past 30 days	9%	6%	3%	9%	9%	11%
Rode with marijuana user, past 30 days	31%	24%	15%	23%	24%	37%
Drive after marijuana use, past 30 days	15%	7%	6%	11%	9%	22%
Carried a weapon on school property, past 30 days	11%	5%	9%	9%	14%	8%
Did not go to school because felt unsafe, past 30 days	8%	3%	5%	3%	6%	8%
Threatened or injured with weapon on school property, past 12 months	7%	4%	7%	7%	10%	9%
In physical fight, past 12 months	30%	22%	24%	33%	34%	32%
In physical fight at school, past 12 months	14%	12%	13%	17%	16%	13%

Personal Safety - Females

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Always/almost always wear seat belt when riding in car	75%	73%	75%	81%	76%	76%
Rode with drinking driver, past 30 days	33%	25%	19%	27%	27%	18%
Drove after drinking, past 30 days	10%	8%	3%	10%	7%	7%
Rode with marijuana user, past 30 days	37%	23%	13%	24%	23%	32%
Drive after marijuana use, past 30 days	17%	9%	5%	10%	7%	15%
Carried a weapon on school property, past 30 days	9%	3%	5%	6%	10%	7%
Did not go to school because felt unsafe, past 30 days	14%	4%	6%	5%	6%	11%
Threatened or injured with weapon on school property, past 12 months	5%	3%	6%	5%	10%	11%
In physical fight, past 12 months	24%	19%	20%	24%	21%	26%
In physical fight at school, past 12 months	16%	6%	9%	11%	5%	9%

Personal Safety - Males

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Always/almost always wear seat belt when riding in car	65%	69%	80%	80%	67%	76%
Rode with drinking driver, past 30 days	29%	15%	15%	21%	17%	26%
Drove after drinking, past 30 days	8%	5%	3%	8%	10%	15%
Rode with marijuana user, past 30 days	25%	26%	17%	23%	26%	43%
Drive after marijuana use, past 30 days	12%	6%	7%	12%	12%	28%
Carried a weapon on school property, past 30 days	11%	8%	14%	11%	18%	9%
Did not go to school because felt unsafe, past 30 days	3%	2%	3%	1%	6%	5%
Threatened or injured with weapon on school property, past 12 months	9%	4%	8%	9%	9%	8%
In physical fight, past 12 months	36%	25%	29%	41%	45%	39%
In physical fight at school, past 12 months	13%	18%	17%	24%	27%	16%

Personal Safety - All

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Were bullied, past 30 days	#	#	#	14%	20%	19%
Bullied someone, past 30 days	#	#	#	28%	29%	25%
Electronically bullied, past 12 months	#	#	#	#	#	22%
Hit by boy/girlfriend on purpose, past 12 months	8%	8%	7%	10%	8%	9%
Purposefully hurt self without wanting to die, ever	#	#	#	#	20%	20%
Planned suicide, past 12 months	22%	14%	15%	17%	11%	14%
Attempted suicide, past 12 months	15%	9%	7%	8%	9%	7%

Personal Safety - Females

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Were bullied, past 30 days	#	#	#	12%	22%	28%
Bullied someone, past 30 days	#	#	#	23%	26%	24%
Electronically bullied, past 12 months	#	#	#	#	#	28%
Hit by boy/girlfriend on purpose, past 12 months	11%	8%	7%	11%	5%	7%
Purposefully hurt self without wanting to die, ever	#	#	#	#	25%	25%
Planned suicide, past 12 months	40%	22%	21%	22%	12%	14%
Attempted suicide, past 12 months	27%	15%	10%	11%	9%	4%

Personal Safety - Males

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Were bullied, past 30 days	#	#	#	15%	18%	11%
Bullied someone, past 30 days	#	#	#	33%	32%	26%
Electronically bullied, past 12 months	#	#	#	#	#	17%
Hit by boy/girlfriend on purpose, past 12 months	4%	7%	6%	10%	10%	10%
Purposefully hurt self without wanting to die, ever	#	#	#	#	16%	16%
Planned suicide, past 12 months	2%	5%	9%	12%	11%	14%
Attempted suicide, past 12 months	2%	3%	4%	6%	10%	9%

Alcohol - All

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Consumed alcohol, ever	81%	77%	69%	66%	69%	71%
Drank alcohol prior to age 13	39%	31%	24%	19%	24%	17%
Consumed alcohol, past 30 days	52%	38%	25%	28%	38%	46%
Binged on alcohol, past 30 days	37%	25%	12%	19%	18%	27%
3 or more days drinking, past 30 days	32%	19%	11%	13%	17%	25%
10 or more days drinking, past 30 days	10%	4%	5%	6%	5%	7%
3 or more days bingeing, past 30 days	17%	8%	6%	8%	10%	13%
Drank on school property, past 30 days	3%	3%	3%	4%	4%	2%

Alcohol - Females

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Consumed alcohol, ever	77%	75%	73%	67%	72%	69%
Drank alcohol prior to age 13	38%	28%	22%	16%	22%	16%
Consumed alcohol, past 30 days	50%	33%	30%	34%	41%	44%
Binged on alcohol, past 30 days	34%	21%	14%	22%	17%	26%
3 or more days drinking, past 30 days	32%	19%	10%	12%	14%	19%
10 or more days drinking, past 30 days	9%	5%	6%	5%	3%	7%
3 or more days bingeing, past 30 days	16%	5%	4%	5%	9%	11%
Drank on school property, past 30 days	5%	5%	5%	5%	2%	1%

Alcohol - Males

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Consumed alcohol, ever	84%	79%	66%	64%	65%	72%
Drank alcohol prior to age 13	41%	34%	25%	22%	26%	18%
Consumed alcohol, past 30 days	53%	44%	21%	24%	34%	47%
Binged on alcohol, past 30 days	40%	31%	10%	17%	19%	27%
3 or more days drinking, past 30 days	32%	20%	13%	14%	18%	31%
10 or more days drinking, past 30 days	11%	3%	5%	7%	7%	7%
3 or more days bingeing, past 30 days	20%	12%	9%	10%	12%	15%
Drank on school property, past 30 days	1%	2%	1%	3%	6%	3%

Alcohol - All

Of students who drank in the past 30 days

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Usually consumed liquor such as vodka, rum, scotch, bourbon, or whiskey	#	#	#	#	47%	51%
Usually consumed beer	#	#	#	#	17%	22%
Usually consumed alcopops	#	#	#	#	29%	21%
Usually consumed wine or wine coolers	#	#	#	#	8%	6%
Someone gave alcohol to me	#	#	#	#	#	45%
Gave someone money to buy alcohol for me	#	#	#	#	#	33%
Got or stole alcohol from home	#	#	#	#	#	13%
Bought in a store	#	#	#	#	#	6%
Stole alcohol from a store or restaurant	#	#	#	#	#	3%

Alcohol - Females

Of students who drank in the past 30 days

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Usually consumed liquor such as vodka, rum, scotch, bourbon, or whiskey	#	#	#	#	47%	45%
Usually consumed beer	#	#	#	#	16%	12%
Usually consumed alcopops	#	#	#	#	32%	33%
Usually consumed wine or wine coolers	#	#	#	#	5%	9%
Someone gave alcohol to me	#	#	#	#	#	46%
Gave someone money to buy alcohol for me	#	#	#	#	#	34%
Got or stole alcohol from home	#	#	#	#	#	11%
Bought in a store	#	#	#	#	#	6%
Stole alcohol from a store or restaurant	#	#	#	#	#	3%

Alcohol - Males

Of students who drank in the past 30 days

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Usually consumed liquor such as vodka, rum, scotch, bourbon, or whiskey	#	#	#	#	46%	55%
Usually consumed beer	#	#	#	#	19%	32%
Usually consumed alcopops	#	#	#	#	23%	11%
Usually consumed wine or wine coolers	#	#	#	#	12%	3%
Someone gave alcohol to me	#	#	#	#	#	43%
Gave someone money to buy alcohol for me	#	#	#	#	#	33%
Got or stole alcohol from home	#	#	#	#	#	14%
Bought in a store	#	#	#	#	#	7%
Stole alcohol from a store or restaurant	#	#	#	#	#	2%

Tobacco - All

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Smoked a cigarette, ever	63%	44%	33%	34%	34%	38%
Smoked cigarettes, past 30 days	30%	15%	10%	19%	22%	25%
Smoked daily, past 30 days	18%	9%	5%	9%	8%	9%
Smoked first cigarette prior to age 13	37%	23%	15%	17%	18%	12%
Smoked 11 or more cigarettes on days smoked, past 30 days	8%	2%	1%	5%	5%	2%
Smoked 20 or more cigarettes on days smoked, past 30 days	1%	0%	0%	1%	4%	0%
Used chewing tobacco, past 30 days	15%	5%	6%	18%	9%	16%
Smoked cigars, past 30 days	14%	9%	10%	#	#	22%

Tobacco - Females

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Smoked a cigarette, ever	66%	50%	34%	33%	30%	41%
Smoked cigarettes, past 30 days	38%	20%	9%	18%	19%	28%
Smoked daily, past 30 days	21%	12%	5%	8%	5%	11%
Smoked first cigarette prior to age 13	37%	27%	13%	15%	16%	15%
Smoked 11 or more cigarettes on days smoked, past 30 days	9%	3%	1%	3%	4%	1%
Smoked 20 or more cigarettes on days smoked, past 30 days	1%	1%	0%	1%	3%	0%
Used chewing tobacco, past 30 days	5%	2%	2%	12%	4%	10%
Smoked cigars, past 30 days	11%	3%	6%	#	#	10%

Tobacco - Males

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Smoked a cigarette, ever	60%	38%	31%	35%	36%	34%
Smoked cigarettes, past 30 days	23%	11%	10%	21%	25%	23%
Smoked daily, past 30 days	15%	6%	5%	10%	10%	7%
Smoked first cigarette prior to age 13	38%	19%	17%	18%	20%	10%
Smoked 11 or more cigarettes on days smoked, past 30 days	8%	1%	2%	7%	6%	2%
Smoked 20 or more cigarettes on days smoked, past 30 days	1%	0%	1%	2%	4%	0%
Used chewing tobacco, past 30 days	25%	8%	11%	23%	15%	22%
Smoked cigars, past 30 days	16%	15%	14%	#	#	34%

Other Drug Use - All

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Smoked marijuana, ever	56%	47%	42%	40%	43%	57%
Tried marijuana prior to age 13	18%	12%	5%	12%	12%	8%
Used marijuana, past 30 days	32%	22%	13%	23%	23%	40%
Used marijuana 10 or more times, past 30 days	13%	7%	7%	9%	15%	25%
Used marijuana on school property, past 30 days	5%	5%	3%	6%	6%	8%
Used cocaine, past 30 days	2%	3%	0%	5%	6%	5%
Use inhalants, ever	27%	16%	11%	9%	19%	13%
Used heroin, ever	3%	2%	2%	4%	4%	4%
Used methamphetamines, ever	12%	6%	4%	8%	7%	6%
Used hallucinogens, ever	19%	10%	6%	13%	11%	15%
Were offered, sold, or given drugs on school property, past 12 months	29%	19%	17%	15%	22%	29%

Other Drug Use - Females

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Smoked marijuana, ever	57%	47%	46%	35%	34%	51%
Tried marijuana prior to age 13	22%	15%	6%	10%	10%	6%
Used marijuana, past 30 days	37%	21%	11%	19%	20%	31%
Used marijuana 10 or more times, past 30 days	16%	8%	6%	9%	6%	20%
Used marijuana on school property, past 30 days	6%	3%	4%	6%	1%	4%
Used cocaine, past 30 days	4%	6%	1%	4%	3%	6%
Use inhalants, ever	32%	18%	14%	9%	21%	14%
Used heroin, ever	4%	3%	3%	4%	1%	4%
Used methamphetamines, ever	11%	9%	4%	7%	4%	5%
Used hallucinogens, ever	24%	15%	6%	15%	7%	12%
Were offered, sold, or given drugs on school property, past 12 months	27%	21%	11%	15%	19%	22%

Other Drug Use - Males

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Smoked marijuana, ever	54%	46%	37%	43%	53%	62%
Tried marijuana prior to age 13	15%	9%	5%	13%	13%	10%
Used marijuana, past 30 days	27%	23%	15%	27%	27%	49%
Used marijuana 10 or more times, past 30 days	11%	6%	8%	10%	24%	29%
Used marijuana on school property, past 30 days	4%	6%	3%	6%	11%	11%
Used cocaine, past 30 days	0%	1%	0%	6%	9%	3%
Use inhalants, ever	23%	14%	8%	10%	17%	13%
Used heroin, ever	1%	0%	2%	4%	7%	3%
Used methamphetamines, ever	12%	4%	4%	8%	9%	7%
Used hallucinogens, ever	15%	4%	6%	11%	15%	17%
Were offered, sold, or given drugs on school property, past 12 months	30%	18%	24%	16%	26%	36%

Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use - All

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Parents would think it is wrong/very wrong for you to smoke	82%	88%	91%	90%	90%	89%
Parents would think it is wrong/very wrong for you to drink	72%	73%	80%	80%	79%	82%
Parents would think it is wrong/very wrong for you to smoke marijuana	82%	90%	92%	87%	84%	81%
Wrong/very wrong for someone your age to smoke	52%	67%	79%	75%	68%	61%
Wrong/very wrong for someone your age to drink alcohol	41%	49%	56%	52%	43%	40%
Wrong/very wrong for someone your age to smoke marijuana	59%	62%	72%	65%	57%	46%
Easy/very easy to obtain cigarettes	76%	76%	67%	76%	75%	77%
Easy/very easy to obtain alcohol	74%	71%	72%	73%	64%	70%
Easy/very easy to obtain marijuana	63%	62%	56%	53%	69%	76%

Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use - Females

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Parents would think it is wrong/very wrong for you to smoke	83%	84%	90%	88%	92%	89%
Parents would think it is wrong/very wrong for you to drink	78%	73%	75%	72%	82%	83%
Parents would think it is wrong/very wrong for you to smoke marijuana	82%	91%	90%	82%	87%	85%
Wrong/very wrong for someone your age to smoke	50%	59%	75%	71%	70%	60%
Wrong/very wrong for someone your age to drink alcohol	45%	49%	52%	43%	43%	40%
Wrong/very wrong for someone your age to smoke marijuana	59%	66%	70%	62%	62%	52%
Easy/very easy to obtain cigarettes	77%	76%	70%	76%	70%	70%
Easy/very easy to obtain alcohol	70%	70%	74%	79%	68%	68%
Easy/very easy to obtain marijuana	67%	62%	61%	53%	68%	66%

Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use - Males

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Parents would think it is wrong/very wrong for you to smoke	80%	91%	92%	91%	88%	90%
Parents would think it is wrong/very wrong for you to drink	66%	72%	86%	87%	75%	80%
Parents would think it is wrong/very wrong for you to smoke marijuana	82%	90%	93%	91%	81%	76%
Wrong/very wrong for someone your age to smoke	53%	74%	83%	79%	66%	63%
Wrong/very wrong for someone your age to drink alcohol	36%	50%	61%	60%	44%	40%
Wrong/very wrong for someone your age to smoke marijuana	59%	57%	73%	67%	54%	40%
Easy/very easy to obtain cigarettes	75%	76%	64%	75%	80%	84%
Easy/very easy to obtain alcohol	78%	73%	70%	68%	60%	72%
Easy/very easy to obtain marijuana	59%	64%	51%	52%	69%	86%

Sexual Behavior and Orientation - All

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Had sexual intercourse, ever	45%	39%	43%	45%	47%	51%
Had sex, past 3 months	33%	29%	30%	39%	36%	#
Had sex before age 13	6%	5%	4%	8%	9%	3%
4 or more sexual partners in lifetime	11%	11%	8%	13%	17%	16%
3 or more sexual partners in last 3 months	2%	4%	2%	5%	7%	#
Used alcohol or drugs prior to last sex (among sexually active)	24%	21%	16%	19%	22%	27%
Used condom at last sex (among sexually active)	60%	58%	55%	61%	63%	61%
Heterosexual	#	#	#	88%	88%	88%
Gay or lesbian, bisexual, not sure	#	#	#	12%	12%	12%

Sexual Behavior and Orientation - Females

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Had sexual intercourse, ever	44%	41%	51%	45%	47%	50%
Had sex, past 3 months	31%	33%	40%	42%	40%	#
Had sex before age 13	6%	4%	3%	6%	8%	3%
4 or more sexual partners in lifetime	12%	11%	9%	14%	17%	18%
3 or more sexual partners in last 3 months	0%	4%	2%	5%	5%	#
Used alcohol or drugs prior to last sex (among sexually active)	25%	21%	16%	18%	20%	30%
Used condom at last sex (among sexually active)	46%	55%	56%	51%	61%	63%
Heterosexual	#	#	#	84%	85%	84%
Gay or lesbian, bisexual, not sure	#	#	#	16%	15%	16%

Sexual Behavior and Orientation - Males

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Had sexual intercourse, ever	47%	38%	36%	45%	46%	52%
Had sex, past 3 months	34%	25%	20%	36%	33%	#
Had sex before age 13	7%	6%	6%	10%	8%	4%
4 or more sexual partners in lifetime	10%	11%	7%	12%	17%	15%
3 or more sexual partners in last 3 months	3%	4%	3%	5%	9%	#
Used alcohol or drugs prior to last sex (among sexually active)	24%	21%	16%	21%	24%	23%
Used condom at last sex (among sexually active)	71%	61%	54%	69%	63%	59%
Heterosexual	#	#	#	93%	91%	93%
Gay or lesbian, bisexual, not sure	#	#	#	7%	9%	7%

Body Image - All

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
BMI between 85th-95th percentile	14%	10%	11%	14%	13%	13%
BMI >95th percentile	9%	11%	6%	12%	12%	16%
Underweight	19%	17%	16%	14%	11%	9%
Right weight	47%	54%	61%	58%	60%	58%
Overweight	34%	29%	23%	28%	28%	33%
Trying to lose weight	48%	42%	40%	40%	45%	42%
Trying to gain weight	21%	14%	16%	16%	18%	14%
Trying to stay at current weight	11%	19%	16%	20%	19%	18%
Doing nothing about weight	19%	26%	28%	24%	19%	25%

Body Image - Females

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
BMI between 85th-95th percentile	8%	13%	9%	13%	13%	16%
BMI >95th percentile	11%	16%	2%	11%	10%	14%
Underweight	8%	13%	9%	6%	9%	2%
Right weight	42%	50%	62%	55%	57%	59%
Overweight	49%	37%	28%	39%	33%	39%
Trying to lose weight	71%	61%	55%	58%	60%	63%
Trying to gain weight	3%	3%	3%	5%	6%	2%
Trying to stay at current weight	12%	16%	18%	13%	19%	18%
Doing nothing about weight	15%	19%	24%	24%	15%	16%

Body Image - Males

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
BMI between 85th-95th percentile	19%	7%	14%	15%	12%	9%
BMI >95th percentile	7%	7%	10%	13%	14%	19%
Underweight	28%	21%	23%	20%	14%	16%
Right weight	54%	58%	59%	61%	63%	57%
Overweight	18%	21%	18%	19%	23%	26%
Trying to lose weight	26%	23%	24%	23%	29%	22%
Trying to gain weight	39%	24%	30%	26%	29%	25%
Trying to stay at current weight	11%	21%	13%	27%	19%	18%
Doing nothing about weight	25%	32%	32%	24%	22%	34%

Nutrition - All

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
2 or more servings of fruit per day, past week	44%	50%	29%	33%	29%	35%
3 or more servings of vegetables per day, past week	9%	17%	10%	14%	11%	20%
5 or more servings of fruit/vegetables per day, past week	23%	24%	17%	21%	19%	26%
Breakfast on 7 or more days, past week	#	#	#	#	38%	38%
Breakfast on 3 or more days, past week	#	#	#	#	64%	76%
Breakfast on 0 days, past week	#	#	#	#	15%	7%
1 or more servings of soda per day, past week	#	#	#	#	27%	26%

Nutrition - Females

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
2 or more servings of fruit per day, past week	40%	50%	33%	33%	32%	38%
3 or more servings of vegetables per day, past week	7%	22%	10%	14%	13%	19%
5 or more servings of fruit/vegetables per day, past week	18%	27%	19%	18%	20%	27%
Breakfast on 3 or more days, past week	#	#	#	#	66%	73%
Breakfast on 7 or more days, past week	#	#	#	#	37%	38%
1 or more servings of soda per day, past week	#	#	#	#	19%	18%

Nutrition - Males

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
2 or more servings of fruit per day, past week	48%	51%	26%	34%	27%	32%
3 or more servings of vegetables per day, past week	12%	13%	9%	14%	9%	19%
5 or more servings of fruit/vegetables per day, past week	27%	21%	16%	24%	19%	24%
Breakfast on 3 or more days, past week	#	#	#	#	63%	78%
Breakfast on 7 or more days, past week	#	#	#	#	38%	36%
1 or more servings of soda per day, past week	#	#	#	#	37%	34%

Physical Activity - All

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
60+ minutes of physical activity every day, past 7 days	#	#	#	#	24%	27%
60+ minutes of physical activity on zero days, past 7 days	#	#	#	#	15%	13%
P.E. at least once per week	30%	49%	42%	53%	45%	49%
P.E. five days per week	22%	44%	41%	49%	39%	42%
Watched tv/used computer for fun, 3 or more hours per day	#	28%	30%	43%	40%	41%
Watched tv/used computer for fun, 5 or more hours per day	#	5%	7%	11%	11%	10%

Physical Activity - Females

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
60+ minutes of physical activity every day, past 7 days	#	#	#	#	18%	20%
60+ minutes of physical activity on zero days, past 7 days	#	#	#	#	16%	21%
P.E. at least once per week	31%	47%	37%	49%	44%	52%
P.E. five days per week	23%	41%	36%	48%	38%	45%
Watched tv/used computer for fun, 3 or more hours per day	#	18%	25%	39%	34%	40%
Watched tv/used computer for fun, 5 or more hours per day	#	3%	6%	8%	8%	12%

Physical Activity - Males

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
60+ minutes of physical activity every day, past 7 days	#	#	#	#	29%	34%
60+ minutes of physical activity on zero days, past 7 days	#	#	#	#	14%	6%
P.E. at least once per week	30%	50%	48%	57%	46%	46%
P.E. five days per week	22%	47%	47%	50%	41%	39%
Watched tv/used computer for fun, 3 or more hours per day	#	38%	36%	47%	45%	42%
Watched tv/used computer for fun, 5 or more hours per day	#	7%	8%	13%	14%	7%

Youth Assets - All

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Earned mostly As	#	33%	32%	32%	30%	22%
Earned mostly Bs	#	39%	40%	44%	37%	40%
Earned mostly Cs	#	20%	15%	15%	20%	26%
Earned mostly Ds or Fs	#	6%	9%	3%	9%	8%
Not sure/not graded	#	3%	3%	6%	4%	4%
Parents talk to you about school at least once a week	70%	75%	71%	73%	79%	76%
Volunteer, 1 or more hours per week	#	55%	46%	39%	41%	41%
Volunteer, 3 or more hours per week	#	17%	13%	9%	8%	11%
Agree/strongly agree that students help decide what goes on at school	#	46%	47%	50%	33%	47%
Agree/strongly agree that you matter to your community	#	43%	39%	40%	39%	40%

Youth Assets - Females

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Earned mostly As	#	38%	35%	37%	33%	28%
Earned mostly Bs	#	39%	41%	44%	36%	40%
Earned mostly Cs	#	18%	16%	12%	19%	22%
Earned mostly Ds or Fs	#	3%	6%	3%	9%	7%
Not sure/not graded	#	3%	3%	5%	3%	2%
Parents talk to you about school at least once a week	74%	71%	71%	75%	77%	74%
Volunteer, 1 or more hours per week	#	53%	51%	31%	42%	37%
Volunteer, 3 or more hours per week	#	16%	18%	9%	7%	12%
Agree/strongly agree that students help decide what goes on at school	#	42%	45%	43%	28%	49%
Agree/strongly agree that you matter to your community	#	33%	33%	37%	29%	37%

Youth Assets - Males

This table includes only the variables on the 2011 YRBS. # = question was not asked that year.

	Year					
	1999	2001	2003	2005	2007	2009
Earned mostly As	#	28%	30%	27%	26%	16%
Earned mostly Bs	#	38%	39%	44%	40%	39%
Earned mostly Cs	#	22%	15%	18%	22%	31%
Earned mostly Ds or Fs	#	10%	13%	4%	8%	9%
Not sure/not graded	#	3%	3%	7%	4%	5%
Parents talk to you about school at least once a week	67%	80%	71%	71%	80%	78%
Volunteer, 1 or more hours per week	#	59%	41%	48%	41%	44%
Volunteer, 3 or more hours per week	#	18%	9%	9%	8%	10%
Agree/strongly agree that students help decide what goes on at school	#	49%	48%	56%	39%	46%
Agree/strongly agree that you matter to your community	#	55%	46%	42%	48%	43%