



Tips for Teens: The Truth About Methamphetamine

Slang--Speed, Meth, Crystal, Crank, Tweak, Go-fast, Ice, Glass, Uppers, Black Beauties

➔ *Get the Facts...*

Methamphetamine affects your brain. In the short term, meth causes mind and mood changes such as anxiety, euphoria, and depression. Long-term effects can include chronic fatigue, paranoid or delusional thinking, and permanent psychological damage.

Methamphetamine affects your body. Over “amping” on any type of speed is pretty risky. Creating a false sense of energy, these drugs push the body faster and further than it's meant to go. It increases the heart rate, blood pressure, and risk of stroke.

Methamphetamine affects your self-control. Meth is a powerfully addictive drug that can cause aggression and violent or psychotic behavior.¹

Methamphetamine is not what it seems. Even speed drugs are not always safe. Giga-jolts of the well-known stimulants caffeine or ephedrine can cause stroke or cardiac arrest when overused or used by people with a sensitivity to them.

Methamphetamine can kill you. An overdose of meth can result in heart failure. Long-term physical effects such as liver, kidney, and lung damage may also kill you.

➔ *Before You Risk It...*

Know the law. Methamphetamine is illegal in all states and highly dangerous.

Get the facts. The ignitable, corrosive, and toxic nature of the chemicals used to produce meth can cause fires, produce toxic vapors, and damage the environment.

Stay informed. In 2001, methamphetamine use sent more people to the emergency room than use of any other club drug. Over half of these cases involved meth in combination with another drug, such as alcohol, heroin, or cocaine.²

Know the risks. There are a lot of risks associated with using methamphetamine, including:

- Meth can cause a severe “crash” after the effects wear off.
- Meth use can cause irreversible damage to blood vessels in the brain.
- Meth users who inject the drug and share needles are at risk for acquiring HIV/AIDS.

Look around you. Not everyone is using methamphetamine. In 2003, only 3.2 percent of 12th graders reported having used methamphetamine.³

Know the Signs...

How can you tell if a friend is using meth? It may not be easy to tell. But there are signs you can look for. Symptoms of methamphetamine use may include:

- Inability to sleep
- Increased sensitivity to noise
- Nervous physical activity, like scratching
- Irritability, dizziness, or confusion
- Extreme anorexia
- Tremors or even convulsions
- Increased heart rate, blood pressure, and risk of stroke
- Presence of inhaling paraphernalia, such as razor blades, mirrors, and straws
- Presence of injecting paraphernalia, such as syringes, heated spoons, or surgical tubing

What can you do to help a friend who is using meth? Be a real friend. You might even save a life. Encourage your friend to stop or seek professional help. For information and referrals, call the National Clearinghouse for Alcohol and Drug Information at 800-729-6686.

Q&A

Q. Isn't methamphetamine less harmful than crack, cocaine, or heroin?

A. Some users get hooked the first time they snort, smoke, or inject meth. Because it can be made from lethal ingredients like battery acid, drain cleaner, lantern fuel, and antifreeze, there is a greater chance of suffering a heart attack, stroke, or serious brain damage with this drug than with other drugs.

Q. Isn't using methamphetamine like using diet pills?

A. No. Though it is easily attainable, methamphetamine is dangerous and addictive. From 1998 to 1999, deaths due to meth rose 38 percent.⁴ In 2002, meth was involved in 17,696 emergency room visits.⁵

Info

To learn more about methamphetamine or obtain referrals to programs in your community, contact one of the following toll-free numbers:

SAMHSA's National Clearinghouse for Alcohol and Drug Information
800-729-6686
TDD 800-487-4889
línea gratis en español
877-767-8432

Curious about the TV ads of the National Youth Anti-Drug Media Campaign? Check out the Web site at www.freevibe.com or visit the Office of National Drug Control Policy Web site at www.whitehousedrugpolicy.gov.

The bottom line: If you know someone who uses meth, urge him or her to stop or get help. If you use meth--stop! The longer you ignore the real facts, the more chances you take with your life.

It's never too late. Talk to your parents, a doctor, a counselor, a teacher, or another adult you trust.

Do it today!

Footnotes

¹ "NIDA Community Drug Alert Bulletin--Club Drugs." National Institute on Drug Abuse, updated 2004.

² *The DAWN Report: Club Drugs, 2001 Update*. Drug Abuse Warning Network, 2002.

³ *Monitoring the Future: National Results on Adolescent Drug Use: Overview of Key Findings, 2003*. National Institute on Drug Abuse, 2004.

⁴ *Prevention Alert: Meth: What's Cooking in Your Neighborhood?* The Substance Abuse and Mental Health Services Administration, 2001.

⁵ *The DAWN Report: Major Drugs of Abuse in ED Visits, 2002 Update*. Drug Abuse Warning Network, 2004